

Comparison of Several Multivariate Means

Shyh-Kang Jeng

Department of Electrical Engineering/
Graduate Institute of Communication/
Graduate Institute of Networking and
Multimedia

1

Outline

- Introduction
- Comparison of Univariate Means
- Paired Comparisons and a Repeated Measures Design
- Comparing Mean Vectors from Two Populations
- Comparison of Several Univariate Population Mean (One-Way ANOVA)

2

Outline

- Comparing Several Multivariate Population Means (One-Way MANOVA)
- Simultaneous Confidence Intervals for Treatment Effects
- Testing for Equality of Covariance Matrices
- Two-Way ANOVA
- Two-Way Multivariate Analysis of Variance

3

Outline

- Profile Analysis
- ANOVA for Repeated Measures
- Repeated Measures Designs and Growth Curves
- Perspectives and Strategy for Analyzing Multivariate Models

4

Outline

- Introduction
- Comparison of Univariate Means
- Paired Comparisons and a Repeated Measures Design
- Comparing Mean Vectors from Two Populations
- Comparison of Several Univariate Population Mean (One-Way ANOVA)

5

Introduction

- Extend previous ideas to handle problems involving the comparison of several mean vectors

6

Outline

- Introduction
- Comparison of Univariate Means
- Paired Comparisons and a Repeated Measures Design
- Comparing Mean Vectors from Two Populations
- Comparison of Several Univariate Population Mean (One-Way ANOVA)

7

Questions

- What is the paired comparison?
- How to design experiments for paired comparison?
- How to test if the population means of paired groups are different?
- How to compute the confidence interval for the difference of population means of paired groups?

8

Questions

- How to compare population means of two populations without paired experiments?
- In such a case, how to estimate the common variance?

9

Scenarios

- To test if the differences are significant between
 - Teaching using Power Point vs. using chalks and blackboard only
 - Drug vs. placebos
 - Processing speed of MP3 player model I of brand A vs. model G of brand B
 - Performance of students going to cram schools vs. those not


10

Paired Comparisons

- Measurements are recorded under different sets of conditions
- See if the responses differ significantly over these sets
- Two or more treatments can be administered to the same or similar experimental units
- Compare responses to assess the effects of the treatments

11

Experiment Design for Paired Comparisons



12

Single Response (Univariate) Case

$$D_j = X_{j1} - X_{j2}, j = 1, 2, \dots, n$$

$$D_j : N(\delta, \sigma_d^2)$$

$$t = \frac{\bar{D} - \delta}{s_d / \sqrt{n}} : t_{n-1}$$

Reject $H_0 : \delta = 0$ in favor of $H_1 : \delta \neq 0$ if $|t| > t_{n-1}(\alpha/2)$

100(1 - α)% confidence interval for δ

$$\bar{d} - t_{n-1}(\alpha/2) \frac{s_d}{\sqrt{n}} \leq \delta \leq \bar{d} + t_{n-1}(\alpha/2) \frac{s_d}{\sqrt{n}}$$

13

Comparing Means from Two Populations

- Without explicitly controlling for unit-to-unit variability, as in the paired comparison case
- Experimental units are randomly assigned to populations
- Applicable to a more general collection of experimental units

14

Assumptions Concerning the Structure of Data

$X_{11}, X_{12}, \dots, X_{1n_1}$: random sample from univariate population with mean μ_1 and variance σ_1^2

$X_{21}, X_{22}, \dots, X_{2n_2}$: random sample from univariate population with mean μ_2 and variance σ_2^2

$X_{11}, X_{12}, \dots, X_{1n_1}$ are independent of $X_{21}, X_{22}, \dots, X_{2n_2}$

Further assumptions when n_1 and n_2 small :

Both populations are univariate normal

$$\sigma_1^2 = \sigma_2^2$$

15

Pooled Estimate of Population Variance

$$\sum_{j=1}^{n_1} (x_{j1} - \bar{x}_1)(x_{j1} - \bar{x}_1)' \approx (n_1 - 1)\sigma^2$$

$$\sum_{j=1}^{n_2} (x_{j2} - \bar{x}_2)(x_{j2} - \bar{x}_2)' \approx (n_2 - 1)\sigma^2$$

$$s_{pooled}^2 = \frac{\sum_{j=1}^{n_1} (x_{j1} - \bar{x}_1)(x_{j1} - \bar{x}_1)' + \sum_{j=1}^{n_2} (x_{j2} - \bar{x}_2)(x_{j2} - \bar{x}_2)'}{n_1 + n_2 - 2}$$

$$= \frac{n_1 - 1}{n_1 + n_2 - 2} s_1^2 + \frac{n_2 - 1}{n_1 + n_2 - 2} s_2^2$$

16

***t*-Statistics for Comparing Two Populations**

$$\begin{aligned}
 &X_{11}, X_{12}, \dots, X_{1n_1} : N(\mu_1, \sigma^2) \\
 &X_{21}, X_{22}, \dots, X_{2n_2} : N(\mu_2, \sigma^2) \\
 &\bar{X}_1 - \bar{X}_2 = \frac{1}{n_1} X_{11} + \dots + \frac{1}{n_1} X_{1n_1} - \frac{1}{n_2} X_{21} + \dots - \frac{1}{n_2} X_{2n_2} \\
 &\quad : N(\mu_1 - \mu_2, \left(\frac{1}{n_1} + \frac{1}{n_2}\right) \sigma^2) \\
 &\Rightarrow t = (\bar{X}_1 - \bar{X}_2 - (\mu_1 - \mu_2)) / \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2}\right) s_{pooled}^2}
 \end{aligned}$$

17

Test of Hypothesis

Reject $H_0 : \mu_1 - \mu_2 = \delta_0$ in favor of $H_1 : \mu_1 - \mu_2 \neq \delta_0$

$$\text{if } \left| \frac{\bar{x}_1 - \bar{x}_2 - \delta_0}{s_{pooled} \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \right| > t_{n_1+n_2-2} \left(\frac{\alpha}{2} \right)$$

18

Outline

- Introduction
- Comparison of Univariate Means
- Paired Comparisons and a Repeated Measures Design
- Comparing Mean Vectors from Two Populations
- Comparison of Several Univariate Population Mean (One-Way ANOVA)

19

Questions

- How to make paired comparison for multivariate data?
- How to use the contrast matrix to carry out paired comparison for multivariate data?
- What is the repeated measures?
- How to test for equality of treatments in a repeated measures?

20

Example 6.1: Effluent Data from Two Labs

Sample j	Commercial lab		State lab of hygiene	
	x_{1j1} (BOD)	x_{1j2} (SS)	x_{2j1} (BOD)	x_{2j2} (SS)
1	6	27	25	15
2	6	23	28	13
3	18	64	36	22
4	8	44	35	29
5	11	30	15	31
6	34	75	44	64
7	28	26	42	30
8	71	124	54	64
9	43	54	34	56
10	33	30	29	20
11	20	14	39	21

Source: Data courtesy of S. Weber.

21

Multivariate Extension: Notations

X_{1j1} = variable 1 under treatment 1

X_{1j2} = variable 2 under treatment 1

\vdots

X_{1jp} = variable p under treatment 1

X_{2j1} = variable 1 under treatment 2

X_{2j2} = variable 2 under treatment 2

\vdots

X_{2jp} = variable p under treatment 2

22

Result 6.1

$$D_{j1} = X_{1j1} - X_{2j1}$$

$$D_{j2} = X_{1j2} - X_{2j2}$$

$$\vdots$$

$$D_{jp} = X_{1jp} - X_{2jp}$$

$$\mathbf{D}_j = [D_{j1}, D_{j2}, \dots, D_{jp}]$$

$$\mathbf{D}_j : N_p(\boldsymbol{\delta}, \boldsymbol{\Sigma}_d), \quad j = 1, 2, \dots, n$$

$$T^2 = n(\bar{\mathbf{D}} - \boldsymbol{\delta})' \mathbf{S}_d^{-1} (\bar{\mathbf{D}} - \boldsymbol{\delta}) : \frac{(n-1)p}{(n-p)} F_{p, n-p}(\alpha)$$

$$\bar{\mathbf{D}} = \frac{1}{n} \sum_{j=1}^n \mathbf{D}_j, \quad \mathbf{S}_d = \frac{1}{n-1} \sum_{j=1}^n (\mathbf{D}_j - \bar{\mathbf{D}})(\mathbf{D}_j - \bar{\mathbf{D}})'$$

23

Test of Hypotheses and Confidence Regions

$\mathbf{d}'_j = [d_{j1}, d_{j2}, \dots, d_{jp}]$: observed differences

Reject $H_0 : \boldsymbol{\delta} = 0$ in favor of $H_1 : \boldsymbol{\delta} \neq 0$ if

$$T^2 = n \bar{\mathbf{d}}' \mathbf{S}_d^{-1} \bar{\mathbf{d}} > \frac{(n-1)p}{n-p} F_{p, n-p}(\alpha)$$

$$\text{Confidence regions: } (\bar{\mathbf{d}} - \boldsymbol{\delta})' \mathbf{S}_d^{-1} (\bar{\mathbf{d}} - \boldsymbol{\delta}) \leq \frac{(n-1)p}{n-p} F_{p, n-p}(\alpha)$$

$$\delta_i : \bar{d}_i \pm \sqrt{\frac{(n-1)p}{n-p} F_{p, n-p}(\alpha)} \sqrt{\frac{s_{d_i}^2}{n}}, \quad \delta_i : \bar{d}_i \pm t_{n-1} \left(\frac{\alpha}{2p} \right) \sqrt{\frac{s_{d_i}^2}{n}}$$

24

Example 6.1: Check Measurements from Two Labs

$$\bar{\mathbf{d}} = \begin{bmatrix} \bar{d}_1 \\ \bar{d}_2 \end{bmatrix} = \begin{bmatrix} -9.36 \\ 13.27 \end{bmatrix}, \quad \mathbf{S}_d = \begin{bmatrix} 199.26 & 88.38 \\ 88.38 & 418.61 \end{bmatrix}$$

$$T^2 = 11 \begin{bmatrix} -9.36 & 13.27 \end{bmatrix} \begin{bmatrix} 0.0055 & -0.0012 \\ -0.0012 & 0.0026 \end{bmatrix} \begin{bmatrix} -9.36 \\ 13.27 \end{bmatrix}$$

$$= 13.6 > \frac{2 \times 10}{9} F_{2,9}(0.05) = 9.47$$

Reject $H_0 : \delta = 0$

$$\delta_1 : -9.36 \pm \sqrt{9.47} \sqrt{199.26/11} \text{ or } (-22.46, 3.74)$$

$$\delta_2 : 13.27 \pm \sqrt{9.47} \sqrt{418.61/11} \text{ or } (-5.71, 32.25)$$

Both includes zero

25

Alternative View

$$\bar{\mathbf{x}}' = [\bar{x}_{11}, \bar{x}_{12}, \dots, \bar{x}_{1p}, \bar{x}_{21}, \bar{x}_{22}, \dots, \bar{x}_{2p}]$$

$$\mathbf{S} = \begin{bmatrix} \mathbf{S}_{11} & \mathbf{S}_{12} \\ \mathbf{S}_{21} & \mathbf{S}_{22} \end{bmatrix}$$

$$\mathbf{C}_{(p \times 2p)} = \begin{bmatrix} 1 & 0 & \dots & 0 & | & -1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 & | & 0 & -1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & | & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & | & 0 & 0 & \dots & -1 \end{bmatrix}$$

$$\mathbf{d}_j = \mathbf{C}\mathbf{x}_j, \quad \bar{\mathbf{d}} = \mathbf{C}\bar{\mathbf{x}}, \quad \mathbf{S}_d = \mathbf{C}\mathbf{S}\mathbf{C}', \quad T^2 = n\bar{\mathbf{x}}'\mathbf{C}'(\mathbf{C}\mathbf{S}\mathbf{C}')^{-1}\mathbf{C}\bar{\mathbf{x}}$$

26


Repeated Measures Design for Comparing Measurements

- q treatments are compared with respect to a single response variable
- Each subject or experimental unit receives each treatment once over successive periods of time

27

Example 6.2: Treatments in an Anesthetics Experiment

- 19 dogs were initially given the drug pentobarbital followed by four treatments



28

Example 6.2: Sleeping-Dog Data

Dog	Treatment			
	1	2	3	4
1	426	609	556	600
2	253	236	392	395
3	359	433	349	357
4	432	431	522	600
5	405	426	513	513
6	324	438	507	539
7	310	312	410	456
8	326	326	350	504
9	375	447	547	548
10	286	286	403	422
11	349	382	473	497
12	429	410	488	547
13	348	377	447	514
14	412	473	472	446
15	347	326	455	468
16	434	458	637	524
17	364	367	432	469
18	420	395	508	531
19	397	556	645	625

Source: Data courtesy of Dr. J. Atlee.

29

Contrast Matrix

$$\mathbf{X}_j = \begin{bmatrix} X_{j1} \\ X_{j2} \\ \vdots \\ X_{jq} \end{bmatrix}, \quad j=1, 2, \dots, n \quad \boldsymbol{\mu} = E(\mathbf{X}_j)$$

$$\begin{bmatrix} \mu_1 - \mu_2 \\ \mu_1 - \mu_3 \\ \vdots \\ \mu_1 - \mu_q \end{bmatrix} = \begin{bmatrix} 1 & -1 & 0 & \dots & 0 \\ 1 & 0 & -1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & -1 \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_q \end{bmatrix} = \mathbf{C}\boldsymbol{\mu}$$

30

Test for Equality of Treatments in a Repeated Measures Design

$\mathbf{X} : N_q(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, \mathbf{C} : contrast matrix

Test of $H_0 : \mathbf{C}\boldsymbol{\mu} = 0$ vs. $H_1 : \mathbf{C}\boldsymbol{\mu} \neq 0$

Reject H_0 if

$$T^2 = n(\mathbf{C}\bar{\mathbf{x}})'(\mathbf{CSC}')^{-1}\mathbf{C}\bar{\mathbf{x}} > \frac{(n-1)(q-1)}{(n-q+1)} F_{q-1, n-q+1}(\alpha)$$

31

Example 6.2: Contrast Matrix

$(\mu_3 + \mu_4) - (\mu_1 + \mu_2) = (\text{Halothane contrast})$

$(\mu_1 + \mu_3) - (\mu_2 + \mu_4) = (\text{CO}_2 \text{ contrast})$

$(\mu_1 + \mu_4) - (\mu_2 + \mu_3) = (\text{H} - \text{CO}_2 \text{ interaction})$

$$\mathbf{C} = \begin{bmatrix} -1 & -1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

32

Example 6.2: Test of Hypotheses

$$\bar{\mathbf{x}} = \begin{bmatrix} 368.21 \\ 404.63 \\ 479.26 \\ 502.89 \end{bmatrix}, \quad \mathbf{S} = \begin{bmatrix} 2819.29 & & & \\ 3568.42 & 7963.14 & & \\ 2943.49 & 5303.98 & 6851.32 & \\ 2295.35 & 4065.44 & 4499.63 & 4878.99 \end{bmatrix}$$

$$\mathbf{C}\bar{\mathbf{x}} = \begin{bmatrix} 209.31 \\ -60.05 \\ -12.79 \end{bmatrix}, \quad \mathbf{CSC}' = \begin{bmatrix} 9432.32 & 1098.92 & 927.62 \\ 1098.92 & 5195.84 & 914.54 \\ 927.62 & 914.54 & 7557.44 \end{bmatrix}$$

$$T^2 = n(\mathbf{C}\bar{\mathbf{x}})'(\mathbf{CSC}')^{-1}(\mathbf{C}\bar{\mathbf{x}}) = 116$$

$$\frac{(n-1)(q-1)}{(n-q+1)} F_{q-1, n-q+1}(0.05) = 10.94$$

Reject $H_0 : \mathbf{C}\boldsymbol{\mu} = 0$

33

Example 6.2: Simultaneous Confidence Intervals

Contrast of halothane influence

$$(\bar{x}_3 + \bar{x}_4) - (\bar{x}_1 + \bar{x}_2) \pm \sqrt{\frac{18(3)}{16} F_{3,16}(0.05)} \frac{\mathbf{c}'\mathbf{S}\mathbf{c}_1}{19} = 209.31 \pm 73.70$$

CO₂ pressure influence

$$-60.05 \pm \sqrt{10.94} \sqrt{\frac{5195.84}{19}} = -60.5 \pm 54.70$$

H - CO₂ "interaction"

$$-12.79 \pm \sqrt{10.94} \sqrt{\frac{7557.44}{19}} = -12.79 \pm 65.97$$

34

Outline

- Introduction
- Comparison of Univariate Means
- Paired Comparisons and a Repeated Measures Design
- Comparing Mean Vectors from Two Populations
- Comparison of Several Univariate Population Mean (One-Way ANOVA)

35

Questions

- How to compare mean vectors from two populations, not forming paired comparison groups?
- How to pool covariance matrices from two populations?
- How to find simultaneous confidence intervals for comparing mean vectors from two populations?

36

Questions

- What is the multivariate Behrens-Fisher problem and how to solve it?

37

Comparing Mean Vectors from Two Populations

- Populations: Sets of experiment settings
- Without explicitly controlling for unit-to-unit variability, as in the paired comparison case
- Experimental units are randomly assigned to populations
- Applicable to a more general collection of experimental units

38

Assumptions Concerning the Structure of Data

$\mathbf{X}_{11}, \mathbf{X}_{12}, \dots, \mathbf{X}_{1n_1}$: random sample from p – variate population with mean vector $\boldsymbol{\mu}_1$ and covariance $\boldsymbol{\Sigma}_1$
 $\mathbf{X}_{21}, \mathbf{X}_{22}, \dots, \mathbf{X}_{2n_2}$: random sample from p – variate population with mean vector $\boldsymbol{\mu}_2$ and covariance $\boldsymbol{\Sigma}_2$
 $\mathbf{X}_{11}, \mathbf{X}_{12}, \dots, \mathbf{X}_{1n_1}$ are independent of $\mathbf{X}_{21}, \mathbf{X}_{22}, \dots, \mathbf{X}_{2n_2}$
 Further assumptions when n_1 and n_2 small :
 Both populations are multivariate normal
 $\boldsymbol{\Sigma}_1 = \boldsymbol{\Sigma}_2$

39

Pooled Estimate of Population Covariance Matrix

$$\begin{aligned} \sum_{j=1}^{n_1} (\mathbf{x}_{j1} - \bar{\mathbf{x}}_1)(\mathbf{x}_{j1} - \bar{\mathbf{x}}_1)' &\approx (n_1 - 1)\boldsymbol{\Sigma} \\ \sum_{j=1}^{n_2} (\mathbf{x}_{j2} - \bar{\mathbf{x}}_2)(\mathbf{x}_{j2} - \bar{\mathbf{x}}_2)' &\approx (n_2 - 1)\boldsymbol{\Sigma} \\ \mathbf{S}_{pooled} &= \frac{\sum_{j=1}^{n_1} (\mathbf{x}_{j1} - \bar{\mathbf{x}}_1)(\mathbf{x}_{j1} - \bar{\mathbf{x}}_1)' + \sum_{j=1}^{n_2} (\mathbf{x}_{j2} - \bar{\mathbf{x}}_2)(\mathbf{x}_{j2} - \bar{\mathbf{x}}_2)'}{n_1 + n_2 - 2} \\ &= \frac{n_1 - 1}{n_1 + n_2 - 2} \mathbf{S}_1 + \frac{n_2 - 1}{n_1 + n_2 - 2} \mathbf{S}_2 \end{aligned}$$

40

Result 6.2

$$\mathbf{X}_{11}, \mathbf{X}_{12}, \dots, \mathbf{X}_{1n_1} : N_p(\boldsymbol{\mu}_1, \boldsymbol{\Sigma})$$

$$\mathbf{X}_{21}, \mathbf{X}_{22}, \dots, \mathbf{X}_{2n_2} : N_p(\boldsymbol{\mu}_2, \boldsymbol{\Sigma})$$

$$\Rightarrow T^2 = [\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)] \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{S}_{pooled} \right]^{-1} [\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)]$$

is distributed as

$$\frac{(n_1 + n_2 - 2)p}{(n_1 + n_2 - p - 1)} F_{p, n_1 + n_2 - p - 1}$$

41

Proof of Result 6.2

$$\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 = \frac{1}{n_1} \mathbf{X}_{11} + \dots + \frac{1}{n_1} \mathbf{X}_{1n_1} - \frac{1}{n_2} \mathbf{X}_{21} + \dots - \frac{1}{n_2} \mathbf{X}_{2n_2}$$

$$: N_p(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2, \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \boldsymbol{\Sigma})$$

$$(n_1 - 1) \mathbf{S}_1 : W_{n_1 - 1}(\boldsymbol{\Sigma}), \quad (n_2 - 1) \mathbf{S}_2 : W_{n_2 - 1}(\boldsymbol{\Sigma})$$

$$(n_1 - 1) \mathbf{S}_1 + (n_2 - 1) \mathbf{S}_2 : W_{n_1 + n_2 - 2}(\boldsymbol{\Sigma})$$

$$T^2 = \left(\frac{1}{n_1} + \frac{1}{n_2} \right)^{-1/2} [\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)] \mathbf{S}_{pooled}^{-1}$$

$$\left(\frac{1}{n_1} + \frac{1}{n_2} \right)^{-1/2} [\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)]$$

$$= N_p(\mathbf{0}, \boldsymbol{\Sigma})' \left[\frac{W_{n_1 + n_2 - 2}(\boldsymbol{\Sigma})}{n_1 + n_2 - 2} \right]^{-1} N_p(\mathbf{0}, \boldsymbol{\Sigma}) : \frac{(n_1 + n_2 - 2)p}{(n_1 + n_2 - p - 1)} F_{p, n_1 + n_2 - p - 1}$$

42

Wishart Distribution

$$w_{n-1}(\mathbf{A} | \boldsymbol{\Sigma}) = \frac{|\mathbf{A}|^{(n-p-2)/2} e^{-\text{tr}[\mathbf{A}\boldsymbol{\Sigma}^{-1}]/2}}{2^{p(n-1)/2} \pi^{p(p-1)/4} |\boldsymbol{\Sigma}|^{(n-1)/2} \prod_{i=1}^p \Gamma\left(\frac{1}{2}(n-i)\right)}$$

\mathbf{A} : positive definite

Properties :

$$\mathbf{A}_1 : W_{m_1}(\mathbf{A}_1 | \boldsymbol{\Sigma}), \quad \mathbf{A}_2 : W_{m_2}(\mathbf{A}_2 | \boldsymbol{\Sigma}) \Rightarrow$$

$$\mathbf{A}_1 + \mathbf{A}_2 : W_{m_1 + m_2}(\mathbf{A}_1 + \mathbf{A}_2 | \boldsymbol{\Sigma})$$

$$\mathbf{A} : W_m(\mathbf{A} | \boldsymbol{\Sigma}) \Rightarrow \mathbf{CAC}' : W_m(\mathbf{CAC}' | \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}')$$

43

Test of Hypothesis

Reject $H_0 : \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2 = \boldsymbol{\delta}_0$ in favor of $H_1 : \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2 \neq \boldsymbol{\delta}_0$

$$\text{if } T^2 = (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - \boldsymbol{\delta}_0)' \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{S}_{pooled} \right]^{-1} (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - \boldsymbol{\delta}_0)$$

$$> \frac{(n_1 + n_2 - 2)p}{n_1 + n_2 - p - 1} F_{p, n_1 + n_2 - p - 1}(\alpha)$$

Note $E(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2) = \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2$

$$\text{Cov}(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2)$$

$$= \text{Cov}(\bar{\mathbf{X}}_1) - \text{Cov}(\bar{\mathbf{X}}_1, \bar{\mathbf{X}}_2) - \text{Cov}(\bar{\mathbf{X}}_2, \bar{\mathbf{X}}_1) + \text{Cov}(\bar{\mathbf{X}}_2)$$

$$= \text{Cov}(\bar{\mathbf{X}}_1) + \text{Cov}(\bar{\mathbf{X}}_2) = \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \boldsymbol{\Sigma}$$

44

Example 6.3: Comparison of Soaps Manufactured in Two Ways

$$n_1 = n_2 = 50$$

$$\bar{\mathbf{x}}_1 = \begin{bmatrix} 8.3 \\ 4.1 \end{bmatrix}, \quad \mathbf{S}_1 = \begin{bmatrix} 2 & 1 \\ 1 & 6 \end{bmatrix}, \quad \bar{\mathbf{x}}_2 = \begin{bmatrix} 10.2 \\ 3.9 \end{bmatrix}, \quad \mathbf{S}_2 = \begin{bmatrix} 2 & 1 \\ 1 & 4 \end{bmatrix}$$

$$\mathbf{S}_{pooled} = \frac{49}{98}\mathbf{S}_1 + \frac{49}{98}\mathbf{S}_2 = \begin{bmatrix} 2 & 1 \\ 1 & 5 \end{bmatrix}, \quad \bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2 = \begin{bmatrix} -1.9 \\ 0.2 \end{bmatrix}$$

Eigenvalues and eigenvectors of \mathbf{S}_{pooled} :

$$\lambda_1 = 5.303, \quad \mathbf{e}_1 = [0.290 \quad 0.957]'$$


$$\lambda_2 = 1.697, \quad \mathbf{e}_2 = [0.957 \quad -0.290]'$$

$$\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \frac{(n_1 + n_2 - 2)p}{n_1 + n_2 - p - 1} F_{p, n_1 + n_2 - p - 1}(0.05) = 0.25$$

$$\sqrt{\lambda_1} \sqrt{0.25} = 1.15, \quad \sqrt{\lambda_2} \sqrt{0.25} = 0.65$$

45

Example 6.3



46

Result 6.3: Simultaneous Confidence Intervals

$$c^2 = \frac{(n_1 + n_2 - 2)p}{n_1 + n_2 - p - 1} F_{p, n_1 + n_2 - p - 1}(\alpha)$$

$$\mathbf{a}'(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2) \pm c \sqrt{\mathbf{a}' \left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{S}_{pooled} \mathbf{a}}$$

will cover $\mathbf{a}'(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)$ for all \mathbf{a}

In particular, $\mu_{1i} - \mu_{2i}$ will be covered by

$$(\bar{X}_{1i} - \bar{X}_{2i}) \pm c \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2} \right) s_{ii, pooled}}$$

47

Example 6.4: Electrical Usage of Homeowners with and without ACs

$$\bar{\mathbf{x}}_1 = \begin{bmatrix} 204.4 \\ 556.6 \end{bmatrix}, \quad \mathbf{S}_1 = \begin{bmatrix} 13825.3 & 23823.4 \\ 23823.4 & 73107.4 \end{bmatrix}, \quad n_1 = 45$$

$$\bar{\mathbf{x}}_2 = \begin{bmatrix} 130.0 \\ 355.0 \end{bmatrix}, \quad \mathbf{S}_2 = \begin{bmatrix} 8632.0 & 19616.7 \\ 19616.7 & 55964.5 \end{bmatrix}, \quad n_2 = 55$$

$$\mathbf{S}_{pooled} = \frac{n_1 - 1}{n_1 + n_2 - 2} \mathbf{S}_1 + \frac{n_2 - 1}{n_1 + n_2 - 2} \mathbf{S}_2$$

$$= \begin{bmatrix} 10963.7 & 21505.5 \\ 21505.5 & 63661.3 \end{bmatrix}$$

$$c^2 = \frac{98(2)}{97} F_{2, 97}(0.05) = 6.26$$

48

Example 6.4: Electrical Usage of Homeowners with and without ACs

95% simultaneous confidence intervals

$$\mu_{11} - \mu_{21} : (204.4 - 130.0) \pm \sqrt{6.26} \sqrt{\left(\frac{1}{45} + \frac{1}{55}\right)} 10963.7$$


$$\text{or } 21.7 \leq \mu_{11} - \mu_{21} \leq 127.1$$

$$\mu_{12} - \mu_{22} : (556.6 - 355.0) \pm \sqrt{6.26} \sqrt{\left(\frac{1}{45} + \frac{1}{55}\right)} 63661.3$$

$$\text{or } 74.7 \leq \mu_{12} - \mu_{22} \leq 328.5$$

49

Example 6.4: 95% Confidence Ellipse



50

Bonferroni Simultaneous Confidence Intervals

$$\mu_{1i} - \mu_{2i} : (\bar{x}_1 - \bar{x}_2) \pm t_{n_1+n_2-2} \left(\frac{\alpha}{2p} \right) \sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2} \right) S_{ii, pooled}}$$

51

Result 6.4

$n_1 - p$ and $n_2 - p$ are large

100% confidence ellipsoid for $\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2$:

$$[\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)] \left[\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right]^{-1}$$

$$[\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)] \leq \chi_p^2(\alpha)$$

Simultaneous confidence intervals for $\mathbf{a}'(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)$:

$$\mathbf{a}'(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2) \pm \sqrt{\chi_p^2(\alpha)} \sqrt{\mathbf{a}' \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right) \mathbf{a}}$$

52

Proof of Result 6.4

$$E(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2) = \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2$$

$$\text{Cov}(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2) = \text{Cov}(\bar{\mathbf{X}}_1) + \text{Cov}(\bar{\mathbf{X}}_2) = \frac{1}{n_1} \boldsymbol{\Sigma}_1 + \frac{1}{n_2} \boldsymbol{\Sigma}_2$$

$$\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 : \text{nearly } N_p \left(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2, \frac{1}{n_1} \boldsymbol{\Sigma}_1 + \frac{1}{n_2} \boldsymbol{\Sigma}_2 \right)$$

$$[\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)] \left(\frac{1}{n_1} \boldsymbol{\Sigma}_1 + \frac{1}{n_2} \boldsymbol{\Sigma}_2 \right)^{-1}$$

$$[\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)] : \chi_p^2$$

$$\boldsymbol{\Sigma}_1 \sim \mathbf{S}_1, \quad \boldsymbol{\Sigma}_2 \sim \mathbf{S}_2$$

53

Remark

If $n_1 = n_2 = n$

$$\frac{n-1}{n+n-2} = \frac{1}{2}$$

$$\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 = \frac{1}{n} (\mathbf{S}_1 + \mathbf{S}_2)$$

$$= \frac{(n-1)\mathbf{S}_1 + (n-1)\mathbf{S}_2}{n+n-2} \left(\frac{1}{n} + \frac{1}{n} \right) = \mathbf{S}_{pooled} \left(\frac{1}{n} + \frac{1}{n} \right)$$

54

Example 6.5

Example 6.4 Data

$$\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 = \begin{bmatrix} 464.17 & 886.08 \\ 886.08 & 2642.15 \end{bmatrix}$$

$$\mu_{11} - \mu_{21} : 74.4 \pm \sqrt{5.99} \sqrt{464.17} \text{ or } (21.7, 127.1)$$

$$\mu_{12} - \mu_{22} : 201.6 \pm \sqrt{5.99} \sqrt{2642.15} \text{ or } (75.8, 327.4)$$

$$H_0 : \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2 = \mathbf{0}$$

$$T^2 = [\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2] \left[\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right]^{-1} [\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2] = 15.66 > \chi_2^2(0.05) = 5.99$$

$$\text{Critical linear combination : } \left[\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right]^{-1} [\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2] = \begin{bmatrix} 0.041 \\ 0.063 \end{bmatrix}$$

55

Multivariate Behrens-Fisher Problem

- ✦ Test $H_0: \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2 = \mathbf{0}$
- ✦ Population covariance matrices are unequal
- ✦ Sample sizes are not large
- ✦ Populations are multivariate normal
- ✦ Both sizes are greater than the number of variables

56

Approximation of T^2 Distribution

$$T^2 = (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)) \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right)^{-1} (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2))$$

$$= \frac{vp}{v-p+1} F_{p, v-p+1}$$

$$v = \frac{p + p^2}{\sum_{i=1}^2 \frac{1}{n_i} \left[\text{tr} \left[\left(\frac{1}{n_i} \mathbf{S}_i \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right)^{-1} \right)^2 \right] + \left[\text{tr} \left[\frac{1}{n_i} \mathbf{S}_i \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right)^{-1} \right]^2 \right] \right]}$$

$$\min(n_1, n_2) \leq v \leq n_1 + n_2$$

57

Confidence Region

$$(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)) \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right)^{-1} (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2 - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2))$$

$$\leq \frac{vp}{v-p+1} F_{p, v-p+1}(\alpha)$$

58

Example 6.6

Example 6.4 data

$$\frac{1}{n_1} \mathbf{S}_1 \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right)^{-1} = \begin{bmatrix} 0.776 & -0.060 \\ -0.092 & 0.646 \end{bmatrix}$$

$$\frac{1}{n_2} \mathbf{S}_2 \left(\frac{1}{n_1} \mathbf{S}_1 + \frac{1}{n_2} \mathbf{S}_2 \right)^{-1} = \begin{bmatrix} 0.224 & -0.060 \\ 0.092 & 0.354 \end{bmatrix}$$

$$v = 77.6$$

$$\frac{vp}{v-p+1} F_{p, v-p+1}(0.05) = \frac{155.2}{76.6} \times 3.12 = 6.32$$

$$T^2 = 15.66 > 6.32, \quad H_0 : \boldsymbol{\mu}_1 - \boldsymbol{\mu}_2 = 0 \text{ is rejected}$$

59

Outline

- Introduction
- Comparison of Univariate Means
- Paired Comparisons and a Repeated Measures Design
- Comparing Mean Vectors from Two Populations
- Comparison of Several Univariate Population Mean (One-Way ANOVA)

60

Questions

- ✦ Why paired comparisons are not good ways to compare several population means?
- ✦ How to compute summed squares (between)?
- ✦ How to compute summed squares (within)?
- ✦ How to compute summed squares (total)?

61

Questions

- ✦ How to calculate the degrees of freedom for summed squares (between)?
- ✦ How to calculate the degrees of freedom for summed squares (within)?
- ✦ How to calculate the degrees of freedom for summed squares (total)?

62

Questions

- ✦ How to compute the F value for testing of the null hypothesis?
- ✦ How are the three kinds of summed squares related?
- ✦ How to explain the geometric meaning of the degrees of freedom for a treatment vector?
- ✦ What is an ANOVA table?

63

Scenarios

- ✦ To test if the following statements are plausible
 - Music compressed by four MP3 compressors are with the same quality
 - Three new drugs are all as effective as a placebo
 - Four brands of beer are equally tasty
 - Lectures, group studying, and computer assisted instruction are equally effective for undergraduate students

64

Comparing Four MP3 Compressors

- ✦ Test four brands, A, B, C, D
- ✦ 10 subjects each brand (40 in total) to provide a satisfaction rating on a 10-point scale
- ✦ Assume that the rating to each brand is a normal distribution, but all four distributions are with the same variance

65

Hypotheses

- ✦ Null hypothesis

$$H_0 : \mu_A = \mu_B = \mu_C = \mu_D$$

- ✦ Alternative hypothesis

$$H_1 : \text{Not all the } \mu\text{s are equal}$$

66

Problem of Using a t -Test

- ✦ Must compare two brands at a time
- ✦ There are 6 possible comparisons
- ✦ Each has a 0.05 chance of being significant by chance
- ✦ Overall chance of significant result, even when no difference exist, approaches $1-(0.95)^6 \sim 0.26$

67

Sample Data

Subject	A	B	C	D
1	4	5	7	2
2	4	5	8	1
3	5	6	7	2
4	5	6	9	3
5	6	7	6	3
6	3	6	3	4
7	4	4	2	5
8	4	5	2	4
9	3	6	2	4
10	4	3	3	3
Mean	4.2	5.3	4.9	3.1

Grand mean: 4.375

*Adapted from: G. R. Norman and D. L. Streiner, *Biostatistics*, 3rd ed.⁶⁸

Thinking in Terms of Signals and Noises

• Signals

- Overall difference among the means of the groups
- Sum of all the squared differences between group means and the overall means

• Noises

- Overall variability within the groups
- Sum of all the squared differences between individual data and their group means

69

Sum of Squares (Between)

$$SS(\text{between}) = n \sum (\bar{x}_\ell - \bar{x})^2$$

$$SS(\text{between}) = 10[(4.2 - 4.375)^2 + (5.3 - 4.375)^2 + (4.9 - 4.375)^2 + (3.1 - 4.375)^2] \\ = 27.875$$

70

Sum of Squares (Within)

$$SS(\text{within}) = \sum_{\ell} \sum_j (x_{\ell j} - \bar{x}_\ell)^2$$

$$SS(\text{within}) = (4 - 4.2)^2 + (4 - 4.2)^2 + \dots + (4 - 4.2)^2 + \\ (5 - 5.3)^2 + (5 - 5.3)^2 + \dots + (3 - 5.3)^2 + \\ (7 - 4.9)^2 + (8 - 4.9)^2 + \dots + (3 - 4.9)^2 + \\ (2 - 3.1)^2 + (1 - 3.1)^2 + \dots + (3 - 3.1)^2 \\ [40 \text{ terms}] \\ = 101.50$$

71

Sum of Squares (Total)

$$SS(\text{total}) = \sum_{\ell} \sum_j (x_{\ell j} - \bar{x})^2$$

$$x_{\ell j} - \bar{x} = (x_{\ell j} - \bar{x}_\ell) + (\bar{x}_\ell - \bar{x})$$

$$(x_{\ell j} - \bar{x})^2 = (x_{\ell j} - \bar{x}_\ell)^2 + 2(x_{\ell j} - \bar{x}_\ell)(\bar{x}_\ell - \bar{x}) + (\bar{x}_\ell - \bar{x})^2$$

$$\sum_j (x_{\ell j} - \bar{x}_\ell) = 0$$

$$\sum_j (x_{\ell j} - \bar{x})^2 = \sum_j (x_{\ell j} - \bar{x}_\ell)^2 + n(\bar{x}_\ell - \bar{x})^2$$

$$SS(\text{total}) = SS(\text{within}) + SS(\text{between})$$

72

Sum of Squares (Total)

$$\begin{aligned}
 SS(\text{total}) &= (4 - 4.375)^2 + (4 - 4.375)^2 + \dots + (4 - 4.375)^2 + \\
 &\quad (5 - 4.375)^2 + (5 - 4.375)^2 + \dots + (3 - 4.375)^2 + \\
 &\quad (7 - 4.375)^2 + (8 - 4.375)^2 + \dots + (3 - 4.375)^2 + \\
 &\quad (2 - 4.375)^2 + (1 - 4.375)^2 + \dots + (3 - 4.375)^2 \\
 &\quad [40 \text{ terms}] \\
 &= 129.375 = 101.50 + 27.875
 \end{aligned}$$

73

χ^2 Distribution

$$X_1 : N(\mu_1, \sigma_1^2), \quad X_2 : N(\mu_2, \sigma_2^2), \quad \dots,$$

$$X_v : N(\mu_v, \sigma_v^2); \quad Z_i = \frac{X_i - \mu_i}{\sigma_i} : N(0,1)$$

$$\chi^2 = \sum_{i=1}^v \left(\frac{x_i - \mu_i}{\sigma_i} \right)^2, \quad v : \text{degrees of freedom (d.f.)}$$

$$f_v(\chi^2) = \begin{cases} \frac{1}{2^{v/2} \Gamma(v/2)} (\chi^2)^{v/2-1} e^{-\chi^2/2}, & \chi^2 > 0 \\ 0, & \chi^2 \leq 0 \end{cases}$$

(Gamma distribution with $\alpha = v/2$)

74

Distribution of Sum of Squares

$$X : N(\mu, \sigma^2)$$

$$S^2 = \frac{1}{n-1} \sum_{j=1}^n (X_j - \bar{X})^2$$

$$(n-1) \frac{S^2}{\sigma^2} : \chi^2 \text{ distribution with } n-1 \text{ degrees of freedom}$$

[proved by moment generating function, see

P. G. Hoel, *Introduction to Mathematical Statistics*, 5th ed.,
John Wiley & Sons, 1984, p. 281]

75

Distribution of Sum of Squares

$$SS(\text{within}) = \sum_{i=1}^g \sum_{j=1}^n (x_{ij} - \bar{x})^2$$

$$\frac{SS(\text{within})}{\sigma^2} : \chi^2 \text{ distribution with}$$

degree of freedom $df(\text{within}) = gn - 1$

$$SS(\text{between}) = \sum_{i=1}^g (\bar{x}_i - \bar{x})^2$$

$$\bar{x} = \frac{1}{gn} \sum_{i=1}^g \sum_{j=1}^n x_{ij} = \frac{1}{g} \sum_{i=1}^g \left(\frac{1}{n} \sum_{j=1}^n x_{ij} \right) = \frac{1}{g} \sum_{i=1}^g \bar{x}_i$$

$$\frac{SS(\text{between})}{\sigma^2} : \chi^2 \text{ distribution with}$$

degree of freedom $df(\text{between}) = g - 1$

76

F-Distribution

χ_1^2, χ_2^2 : independent, with d.f. f_1 and f_2 , respectively


$$F = \frac{\chi_1^2 / f_1}{\chi_2^2 / f_2}, F > 0$$

$$f(F) = \frac{\Gamma\left(\frac{f_1 + f_2}{2}\right)}{\Gamma\left(\frac{f_1}{2}\right)\Gamma\left(\frac{f_2}{2}\right)} \left(\frac{f_1}{f_2}\right)^{\frac{f_1}{2}} \frac{F^{\frac{f_1}{2}-1}}{\left(1 + \frac{f_1 F}{f_2}\right)^{(f_1 + f_2)/2}}$$

: F_{f_1, f_2}

77

F-Distribution



78

Distribution of F

$$F = \frac{SS(between) / df(between)}{SS(within) / df(within)} :$$

F distribution of degree of freedoms

$df(between)$ and $df(within)$

79

Expected Values of Sum of Squares

if no difference between groups

$$E[SS(between) / df(between)] = \sigma_{err}^2$$

if no difference within groups

$$E[SS(between) / df(between)] = n\sigma_{bet}^2$$

if both differences can happen

$$E[SS(between) / df(between)] = n\sigma_{bet}^2 + \sigma_{err}^2$$

Thus, if H_0 is invalid

$$\frac{E[SS(between) / df(between)]}{E[SS(within) / df(within)]} = \frac{n\sigma_{bet}^2 + \sigma_{err}^2}{\sigma_{err}^2} > 1$$

$F > 1$

80

Degrees of Freedom

$$df(\text{between}) = g - 1 = 4 - 1 = 3$$

$$df(\text{within}) = g(n - 1) = 4(10 - 1) = 36$$

$$\begin{aligned} df(\text{total}) &= gn - 1 = gn - g + g - 1 \\ &= df(\text{within}) + df(\text{between}) \\ &= 40 - 1 = 39 = 36 + 3 \end{aligned}$$

81

ANOVA Summary

Source	Sum of Squares	df	Mean square	F
Between	27.875	3	9.292	3.296
Within	101.500	36	2.819	
Total	129.375	39		

82

Hypothesis Testing

$$F = 3.296 > F_{3,36}(0.05) = 2.86$$

reject $H_0 : \mu_A = \mu_B = \mu_C = \mu_D$
at 0.05 significance level

83

Univariate ANOVA

$X_{\ell 1}, X_{\ell 2}, \dots, X_{\ell n_\ell}$: random sample from $N(\mu_\ell, \sigma^2)$

$\ell = 1, 2, \dots, g$

Null hypothesis $H_0 : \mu_1 = \mu_2 = \dots = \mu_g$

Reparameterization

$$\mu_\ell = \mu + \tau_\ell$$

$$H_0 : \tau_1 = \tau_2 = \dots = \tau_g = 0$$

$$X_{\ell j} = \mu + \tau_\ell + e_{\ell j}, \quad e_{\ell j} : N(0, \sigma^2), \quad \sum_{\ell=1}^g n_\ell \tau_\ell = 0$$

$$x_{\ell j} = \bar{x} + (\bar{x}_\ell - \bar{x}) + (x_{\ell j} - \bar{x}_\ell)$$

84

Univariate ANOVA

$$(x_{ij} - \bar{x})^2 = (\bar{x}_\ell - \bar{x})^2 + (x_{ij} - \bar{x}_\ell)^2 + 2(\bar{x}_\ell - \bar{x})(x_{ij} - \bar{x}_\ell)$$

$$\sum_{j=1}^{n_\ell} (x_{ij} - \bar{x}_\ell) = 0$$

$$\sum_{j=1}^{n_\ell} (x_{ij} - \bar{x})^2 = n_\ell (\bar{x}_\ell - \bar{x})^2 + \sum_{j=1}^{n_\ell} (x_{ij} - \bar{x}_\ell)^2$$

$$\sum_{\ell=1}^g \sum_{j=1}^{n_\ell} (x_{ij} - \bar{x})^2 = \sum_{\ell=1}^g n_\ell (\bar{x}_\ell - \bar{x})^2 + \sum_{\ell=1}^g \sum_{j=1}^{n_\ell} (x_{ij} - \bar{x}_\ell)^2$$

$$(SS_{cor}) = (SS_{tr}) + (SS_{res})$$

$$\sum_{\ell=1}^g \sum_{j=1}^{n_\ell} x_{ij}^2 = (n_1 + n_2 + \dots + n_g) \bar{x}^2 + \sum_{\ell=1}^g n_\ell (\bar{x}_\ell - \bar{x})^2 + \sum_{\ell=1}^g \sum_{j=1}^{n_\ell} (x_{ij} - \bar{x}_\ell)^2$$

$$(SS_{obs}) = (SS_{mean}) + (SS_{tr}) + (SS_{res})$$

85

Concept of Degrees of Freedom

$$\mathbf{y}' = [x_{11}, \dots, x_{1n_1}, x_{21}, \dots, x_{2n_2}, \dots, x_{gn_g}] : \text{d.f. } n = n_1 + n_2 + \dots + n_g$$

Treatment vector

$$\begin{pmatrix} 1 \\ \vdots \\ 1 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + (\bar{x}_1 - \bar{x}) \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots + (\bar{x}_g - \bar{x}) \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 0 \\ 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} = (\bar{x}_1 - \bar{x}) \mathbf{u}_1 + (\bar{x}_2 - \bar{x}) \mathbf{u}_2 + \dots + (\bar{x}_g - \bar{x}) \mathbf{u}_g$$

86

Concept of Degrees of Freedom

$$\mathbf{1} = [1, \dots, 1] = \mathbf{u}_1 + \mathbf{u}_2 + \dots + \mathbf{u}_g$$

Treatment vector and $\mathbf{1}$ are all on the hyperplane spanned by $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_g$: d.f. g

$\mathbf{1}$ is perpendicular to the treatment vector

\therefore mean vector $\bar{x}\mathbf{1}$: d.f. $g - 1$

Residual vector

$$\mathbf{e} = \mathbf{y} - \bar{x}\mathbf{1} = [(x_1 - \bar{x})\mathbf{u}_1 + (x_2 - \bar{x})\mathbf{u}_2 + \dots + (x_g - \bar{x})\mathbf{u}_g]$$

perpendicular to the hyperplane spanned by $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_g$

\therefore d.f. of \mathbf{e} : $n - g$

87

Univariate ANOVA

ANOVA TABLE FOR COMPARING UNIVARIATE POPULATION MEANS

Source of variation	Sum of squares (SS)	Degrees of freedom (d.f.)
Treatments	$SS_{tr} = \sum_{\ell=1}^g n_\ell (\bar{x}_\ell - \bar{x})^2$	$g - 1$
Residual (Error)	$SS_{res} = \sum_{\ell=1}^g \sum_{j=1}^{n_\ell} (x_{ij} - \bar{x}_\ell)^2$	$\sum_{\ell=1}^g n_\ell - g$
Total (corrected for the mean)	$SS_{cor} = \sum_{\ell=1}^g \sum_{j=1}^{n_\ell} (x_{ij} - \bar{x})^2$	$\sum_{\ell=1}^g n_\ell - 1$

88

Univariate ANOVA

Reject $H_0 : \tau_1 = \tau_2 = \dots = \tau_g = 0$ at level α if

$$F = \frac{SS_{tr}/(g-1)}{SS_{res}/\left(\sum_{\ell=1}^g n_{\ell} - g\right)} > F_{g-1, \sum n_{\ell} - g}(\alpha)$$

$$\frac{1}{1 + SS_{tr}/SS_{res}} = \frac{SS_{res}}{SS_{res} + SS_{tr}}$$

89

Examples 6.7 & 6.8

$$\begin{pmatrix} 9 & 6 & 9 \\ 0 & 2 & \\ 3 & 1 & 2 \end{pmatrix} = \begin{pmatrix} 4 & 4 & 4 \\ 4 & 4 & \\ 4 & 4 & 4 \end{pmatrix} + \begin{pmatrix} 4 & 4 & 4 \\ -3 & -3 & \\ -2 & -2 & -2 \end{pmatrix} + \begin{pmatrix} 1 & -2 & 1 \\ -1 & 1 & \\ 1 & -1 & 0 \end{pmatrix}$$

$$SS_{obs} = 216, SS_{mean} = 128$$

$$SS_{tr} = 78, \text{d.f.} = 3 - 1 = 2$$

$$SS_{res} = 10, \text{d.f.} = (3 + 2 + 3) - 3 = 5$$

$$F = \frac{SS_{tr}/(g-1)}{SS_{res}/(\sum n_{\ell} - g)} = \frac{78/2}{10/5} = 19.5 > F_{2,5}(0.01) = 13.27$$

$H_0 : \tau_1 = \tau_2 = \tau_3 = 0$ is rejected at the 1% level

90

Outline

- Comparing Several Multivariate Population Means (One-Way MANOVA)
- Simultaneous Confidence Intervals for Treatment Effects
- Testing for Equality of Covariance Matrices
- Two-Way ANOVA
- Two-Way Multivariate Analysis of Variance

91

Questions

- What is the one-way MANOVA table?
- How to compute Wilk's lambda for MANOVA?
- How to test the equality of several mean vectors from the Wilk's lambda?
- How to test the equality of several mean vectors for large sample size?

92

Questions

- What are other statistics used in statistical software package for one-way MANOVA?

93

Scenario: Example 6.10, Nursing Home Data

- Nursing homes can be classified by the owners: private (271), non-profit (138), government (107)
- Costs: nursing labor, dietary labor, plant operation and maintenance labor, housekeeping and laundry labor
- To investigate the effects of ownership on costs

94

One-Way MANOVA

Population 1: $\mathbf{X}_{11}, \mathbf{X}_{12}, \dots, \mathbf{X}_{1n_1}$

Population 2: $\mathbf{X}_{21}, \mathbf{X}_{22}, \dots, \mathbf{X}_{2n_2}$

\vdots

Population g : $\mathbf{X}_{g1}, \mathbf{X}_{g2}, \dots, \mathbf{X}_{gn_g}$

MANOVA (Multivariate ANalysis Of VAriance)
is used to investigate whether the population mean vectors are the same, and, if not, which mean components differ significantly

95

Assumptions about the Data

$\mathbf{X}_{\ell 1}, \mathbf{X}_{\ell 2}, \dots, \mathbf{X}_{\ell n_\ell}$: random sample from a population
with mean $\boldsymbol{\mu}_\ell$, $\ell = 1, 2, \dots, g$

Random sample from different populations are
independent

All populations have a common covariance
matrix $\boldsymbol{\Sigma}$

Each population is multivariate normal

96

MANOVA

$$\mathbf{X}_{\ell j} = \boldsymbol{\mu} + \boldsymbol{\tau}_{\ell} + \mathbf{e}_{\ell j}; j = 1, 2, \dots, n_{\ell}; \ell = 1, 2, \dots, g$$

$$\mathbf{e}_{\ell j} : N_p(\mathbf{0}, \boldsymbol{\Sigma}), \quad \boldsymbol{\mu} : \text{overall mean (level)}$$

$$\boldsymbol{\tau}_{\ell} : \ell\text{th treatment effect, } \sum_{\ell=1}^g n_{\ell} \boldsymbol{\tau}_{\ell} = \mathbf{0}$$

$$\mathbf{x}_{\ell j} = \bar{\mathbf{x}} + (\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}}) + (\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell}) = \hat{\boldsymbol{\mu}} + \hat{\boldsymbol{\tau}}_{\ell} + \hat{\mathbf{e}}_{\ell j}$$

$$\sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}})' = \sum_{\ell=1}^g n_{\ell} (\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}})'$$

$$+ \sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})' = \mathbf{B} + \mathbf{W}$$

97

MANOVA

MANOVA TABLE FOR COMPARING POPULATION MEAN VECTORS

Source of variation	Matrix of sum of squares and cross products (SSP)	Degrees of freedom (d.f.)
Treatment	$\mathbf{B} = \sum_{\ell=1}^g n_{\ell} (\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}})'$	$g - 1$
Residual (Error)	$\mathbf{W} = \sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})'$	$\sum_{\ell=1}^g n_{\ell} - g$
Total (corrected for the mean)	$\mathbf{B} + \mathbf{W} = \sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}})'$	$\sum_{\ell=1}^g n_{\ell} - 1$

98

MANOVA

$$\mathbf{W} = \sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})'$$

$$= (n_1 - 1)\mathbf{S}_1 + (n_2 - 1)\mathbf{S}_2 + \dots + (n_g - 1)\mathbf{S}_g$$

Reject $H_0 : \boldsymbol{\tau}_1 = \boldsymbol{\tau}_2 = \dots = \boldsymbol{\tau}_g = \mathbf{0}$ if Wilk's lambda

$$\Lambda^* = \frac{|\mathbf{W}|}{|\mathbf{B} + \mathbf{W}|} = \frac{\left| \sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}}_{\ell})' \right|}{\left| \sum_{\ell=1}^g \sum_{j=1}^{n_{\ell}} (\mathbf{x}_{\ell j} - \bar{\mathbf{x}})(\mathbf{x}_{\ell j} - \bar{\mathbf{x}})' \right|}$$

is too small

99

Distribution of Wilk's Lambda

No. of variables	No. of groups	Sampling distribution for multivariate normal data
$p = 1$	$g \geq 2$	$\left(\frac{\sum n_{\ell} - g}{g - 1} \right) \left(\frac{1 - \Lambda^*}{\Lambda^*} \right) \sim F_{g-1, \sum n_{\ell} - g}$
$p = 2$	$g \geq 2$	$\left(\frac{\sum n_{\ell} - g - 1}{g - 1} \right) \left(\frac{1 - \sqrt{\Lambda^*}}{\sqrt{\Lambda^*}} \right) \sim F_{2(g-1), 2(\sum n_{\ell} - g - 1)}$
$p \geq 1$	$g = 2$	$\left(\frac{\sum n_{\ell} - p - 1}{p} \right) \left(\frac{1 - \Lambda^*}{\Lambda^*} \right) \sim F_{p, \sum n_{\ell} - p - 1}$
$p \geq 1$	$g = 3$	$\left(\frac{\sum n_{\ell} - p - 2}{p} \right) \left(\frac{1 - \sqrt{\Lambda^*}}{\sqrt{\Lambda^*}} \right) \sim F_{2p, 2(\sum n_{\ell} - p - 2)}$

100

Test of Hypothesis for Large Size

If H_0 is true and $\sum n_\ell = n$ is large,

$$-\left(n-1-\frac{p+g}{2}\right) \ln \Lambda^* : \chi^2_{p(g-1)}$$

Reject H_0 at significance level α if

$$-\left(n-1-\frac{p+g}{2}\right) \ln \left(\frac{|\mathbf{W}|}{|\mathbf{B} + \mathbf{W}|} \right) > \chi^2_{p(g-1)}(\alpha)$$

101

Popular MANOVA Statistics Used in Statistical Packages

$$\text{Wilk's lambda } \Lambda^* = \frac{|\mathbf{W}|}{|\mathbf{B} + \mathbf{W}|}$$

$$\text{Lawley - Hotelling trace} = \text{tr}[\mathbf{B}\mathbf{W}^{-1}]$$

$$\text{Pillai trace} = \text{tr}[\mathbf{B}(\mathbf{B} + \mathbf{W})^{-1}]$$

Roy's largest root =

maximum eigenvalue of $\mathbf{W}(\mathbf{B} + \mathbf{W})^{-1}$

102

Example 6.9

$$\begin{pmatrix} \begin{bmatrix} 9 \\ 3 \\ 0 \\ 4 \\ 3 \\ 8 \end{bmatrix} & \begin{bmatrix} 6 \\ 2 \\ 2 \\ 0 \\ 1 \\ 9 \end{bmatrix} & \begin{bmatrix} 9 \\ 7 \\ 7 \\ 2 \\ 7 \\ 7 \end{bmatrix} \end{pmatrix} \quad \bar{\mathbf{x}}_1 = \begin{bmatrix} 8 \\ 4 \end{bmatrix}, \bar{\mathbf{x}}_2 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \bar{\mathbf{x}}_3 = \begin{bmatrix} 4 \\ 5 \end{bmatrix}, \bar{\mathbf{x}} = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$$

$$\begin{pmatrix} 9 & 6 & 9 \\ 0 & 2 & \\ 3 & 1 & 2 \end{pmatrix}, SS_{obs} = SS_{mean} + SS_{tr} + SS_{res} = 128 + 78 + 10 = 216$$

$$\begin{pmatrix} 3 & 2 & 7 \\ 4 & 0 & \\ 8 & 9 & 7 \end{pmatrix}, SS_{obs} = SS_{mean} + SS_{tr} + SS_{res} = 200 + 48 + 24 = 272$$

103

Example 6.8

$$\begin{pmatrix} 9 & 6 & 9 \\ 0 & 2 & \\ 3 & 1 & 2 \end{pmatrix} = \begin{pmatrix} 4 & 4 & 4 \\ 4 & 4 & \\ 4 & 4 & 4 \end{pmatrix} + \begin{pmatrix} 4 & 4 & 4 \\ -3 & -3 & \\ -2 & -2 & -2 \end{pmatrix} + \begin{pmatrix} 1 & -2 & 1 \\ -1 & 1 & \\ 1 & -1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 3 & 2 & 7 \\ 4 & 0 & \\ 8 & 9 & 7 \end{pmatrix} = \begin{pmatrix} 5 & 5 & 5 \\ 5 & 5 & \\ 5 & 5 & 5 \end{pmatrix} + \begin{pmatrix} -1 & -1 & -1 \\ -3 & -3 & \\ 3 & 3 & 3 \end{pmatrix} + \begin{pmatrix} -1 & -2 & 3 \\ 2 & -2 & \\ 0 & 1 & -1 \end{pmatrix}$$

Cross products

Mean : $8 \times 4 \times 5 = 160$

Treatment : $3 \times 4 \times (-1) + 2 \times (-3) \times (-3) + 3 \times (-2) \times 3 = -12$

Residual : $1 \times (-1) + (-2) \times (-2) + 1 \times 3 + \dots + 0 \times (-1) = 1$

Total : $9 \times 3 + 6 \times 2 + 9 \times 7 + \dots + 2 \times 7 = 149$

104

Example 6.9

Source of variation	Matrix of sum of squares and cross products	Degrees of freedom
Treatment	$\begin{bmatrix} 78 & -12 \\ -12 & 48 \end{bmatrix}$	$3 - 1 = 2$
Residual	$\begin{bmatrix} 10 & 1 \\ 1 & 24 \end{bmatrix}$	$3 + 2 + 3 - 3 = 5$
Total (corrected)	$\begin{bmatrix} 88 & -11 \\ -11 & 72 \end{bmatrix}$	7

105

Example 6.9

$$\Lambda^* = \frac{|\mathbf{W}|}{|\mathbf{B} + \mathbf{W}|} = \frac{\begin{vmatrix} 10 & 1 \\ 1 & 24 \end{vmatrix}}{\begin{vmatrix} 88 & -11 \\ -11 & 72 \end{vmatrix}} = 0.0385$$

$$\left(\frac{1 - \sqrt{\Lambda^*}}{\sqrt{\Lambda^*}} \right) \frac{\sum n_\ell - g - 1}{g - 1} = \left(\frac{1 - \sqrt{0.0385}}{\sqrt{0.0385}} \right) \frac{8 - 3 - 1}{3 - 1}$$

$$= 8.19 > F_{2(g-1), 2\sum n_\ell - g - 1}(0.01) = F_{4,8}(0.01) = 7.01$$

Reject H_0

106

Example 6.10: Nursing Home Data

- Nursing homes can be classified by the owners: private (271), non-profit (138), government (107)
- Costs: nursing labor, dietary labor, plant operation and maintenance labor, housekeeping and laundry labor
- To investigate the effects of ownership on costs

107

Example 6.10

Group	Number of observations	Sample mean vectors
$\ell = 1$ (private)	$n_1 = 271$	
$\ell = 2$ (nonprofit)	$n_2 = 138$	$\bar{\mathbf{x}}_1 = \begin{bmatrix} 2.066 \\ .480 \\ .082 \\ .360 \end{bmatrix}; \bar{\mathbf{x}}_2 = \begin{bmatrix} 2.167 \\ .596 \\ .124 \\ .418 \end{bmatrix}; \bar{\mathbf{x}}_3 = \begin{bmatrix} 2.273 \\ .521 \\ .125 \\ .383 \end{bmatrix}$
$\ell = 3$ (government)	$n_3 = 107$	
$\sum_{\ell=1}^3 n_\ell = 516$		
Sample covariance matrices		
$\mathbf{S}_1 = \begin{bmatrix} .291 & & & \\ -.001 & .011 & & \\ .002 & .000 & .001 & \\ .010 & .003 & .000 & .010 \end{bmatrix};$	$\mathbf{S}_2 = \begin{bmatrix} .561 & & & \\ .011 & .025 & & \\ .001 & .004 & .005 & \\ .037 & .007 & .002 & .019 \end{bmatrix};$	
$\mathbf{S}_3 = \begin{bmatrix} .261 & & & \\ .030 & .017 & & \\ .003 & -.000 & .004 & \\ .018 & .006 & .001 & .013 \end{bmatrix}$		

Source: Data courtesy of State of Wisconsin Department of Health and Social Services.

108

Example 6.10

$$\mathbf{W} = (n_1 - 1)\mathbf{S}_1 + (n_2 - 1)\mathbf{S}_2 + (n_3 - 1)\mathbf{S}_3$$

$$= \begin{bmatrix} 182.962 & & & \\ 4.408 & 8.200 & & \\ 1.695 & 0.633 & 1.484 & \\ 9.581 & 2.428 & 0.394 & 6.538 \end{bmatrix}$$

$$\bar{\mathbf{x}} = \frac{n_1 \bar{\mathbf{x}}_1 + n_2 \bar{\mathbf{x}}_2 + n_3 \bar{\mathbf{x}}_3}{n_1 + n_2 + n_3} = [2.136 \quad 0.519 \quad 0.102 \quad 0.380]'$$

$$\mathbf{B} = \sum_{\ell=1}^g n_{\ell} (\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell} - \bar{\mathbf{x}})' = \begin{bmatrix} 3.475 & & & \\ 1.111 & 1.225 & & \\ 0.821 & 0.453 & 0.235 & \\ 0.584 & 0.610 & 0.230 & 0.304 \end{bmatrix}$$

109

Example 6.10

$$\Lambda^* = \frac{|W|}{|B+W|} = 0.7714$$

$$\left(\frac{\sum n_{\ell} - p - 2}{p} \right) \left(\frac{1 - \sqrt{\Lambda^*}}{\sqrt{\Lambda^*}} \right) = 17.67$$

$$F_{2 \times 4, 2 \times 510}(0.01) \approx \chi_8^2(0.01)/8 = 2.51$$

or, approximate analysis

$$-(n-1-(p+g)/2) \ln \left(\frac{|W|}{|B+W|} \right) = 132.76$$

$$\chi_{p(g-1)}^2(0.01) = \chi_8^2(0.01) = 20.09$$

Reject H_0 by both analyses

110

Outline

- Comparing Several Multivariate Population Means (One-Way MANOVA)
- Simultaneous Confidence Intervals for Treatment Effects
- Testing for Equality of Covariance Matrices
- Two-Way ANOVA
- Two-Way Multivariate Analysis of Variance

111

Questions

- What are the Bonferroni Intervals for Treatment Effects?

112

Bonferroni Intervals for Treatment Effects

$$\hat{\tau}_{ki} = \bar{x}_{ki} - \bar{x}_i, \quad \hat{\tau}_{ki} - \hat{\tau}_{\ell i} = \bar{x}_{ki} - \bar{x}_{\ell i}$$

$$\text{Var}(\hat{\tau}_{ki} - \hat{\tau}_{\ell i}) = \text{Var}(\bar{x}_{ki} - \bar{x}_{\ell i}) = \left(\frac{1}{n_k} + \frac{1}{n_\ell} \right) \sigma_{ii}$$

$$\mathbf{W} = (n_1 - 1)\mathbf{S}_1 + (n_2 - 1)\mathbf{S}_2 + \cdots + (n_g - 1)\mathbf{S}_g \\ = (n - g)\mathbf{S}_{pooled} \approx (n - g)\mathbf{\Sigma}$$

$$\text{Var}(\hat{\tau}_{ki} - \hat{\tau}_{\ell i}) \approx \left(\frac{1}{n_k} + \frac{1}{n_\ell} \right) \frac{w_{ii}}{(n - g)}$$

$$m = pg(g - 1) / 2$$

113

Result 6.5: Bonferroni Intervals for Treatment Effects

With confidence at least $(1 - \alpha)$

$\tau_{ki} - \tau_{\ell i}$ belongs to

$$\bar{x}_{ki} - \bar{x}_{\ell i} \pm t_{n-g} \left(\frac{\alpha}{pg(g-1)} \right) \sqrt{\frac{w_{ii}}{n-g} \left(\frac{1}{n_k} + \frac{1}{n_\ell} \right)}$$

114

Example 6.11: Example 6.10 Data

$$\hat{\tau}_1 = \bar{x}_1 - \bar{x} = [-0.070 \quad -0.039 \quad -0.020 \quad -0.020]$$

$$\hat{\tau}_3 = \bar{x}_3 - \bar{x} = [0.137 \quad 0.002 \quad 0.023 \quad 0.003]$$

$$\hat{\tau}_{13} - \hat{\tau}_{33} = -0.20 - 0.023 = -0.043, n = 516$$

$$\sqrt{\left(\frac{1}{n_1} + \frac{1}{n_3} \right) \frac{w_{33}}{n-g}} = \sqrt{\left(\frac{1}{271} + \frac{1}{107} \right) \frac{1.484}{516-3}} = 0.00614$$

$$t_{513}(0.05/4 \times 3 \times 2) = 2.87$$

95% simultaneous confidence interval for $\tau_{13} - \tau_{33}$

$$-0.043 \pm 2.87 \times 0.00614 \text{ or } (-0.061, -0.025)$$

95% simultaneous confidence intervals for

$$\tau_{13} - \tau_{23} \text{ and } \tau_{23} - \tau_{33} : (-0.058, -0.026), (-0.021, 0.019)$$

115

Outline

- Comparing Several Multivariate Population Means (One-Way MANOVA)
- Simultaneous Confidence Intervals for Treatment Effects
- Testing for Equality of Covariance Matrices
- Two-Way ANOVA
- Two-Way Multivariate Analysis of Variance

116

Questions

- How to test if the covariance matrices of several populations are equal? (Box's *M*-Test)

117

Test for Equality of Covariance Matrices

- With g populations, null hypothesis

$$H_0: \Sigma_1 = \Sigma_2 = \dots = \Sigma_g = \Sigma$$

- Assume multivariate normal populations

- Likelihood ratio statistic for testing

$$H_0 \quad \Lambda = \prod_{\ell} \left(\frac{|\mathbf{S}_{\ell}|}{|\mathbf{S}_{pooled}|} \right)^{(n_{\ell}-1)/2}$$

$$\mathbf{S}_{pooled} = \frac{1}{\sum_{\ell} (n_{\ell} - 1)} \{ (n_1 - 1)\mathbf{S}_1 + \dots + (n_g - 1)\mathbf{S}_g \}$$

118

Box's *M*-Test

$$M = -2 \ln \Lambda = \left[\sum_{\ell} (n_{\ell} - 1) \right] \ln |\mathbf{S}_{pooled}| - \sum_{\ell} [(n_{\ell} - 1) \ln |\mathbf{S}_{\ell}|]$$

$$u = \left[\sum_{\ell} \frac{1}{(n_{\ell} - 1)} - \frac{1}{\sum_{\ell} (n_{\ell} - 1)} \right] \left[\frac{2p^2 + 3p - 1}{6(p+1)(g-1)} \right]$$

$$C = (1 - u)M : \text{approximate } \chi^2_{\nu}$$

$$\nu = \frac{1}{2} p(p+1)(g-1)$$

$$\text{Reject } H_0 \text{ if } C > \chi^2_{\nu}(\alpha)$$

119

Example 6.12

- Example 6.10 - nursing home data

$$g = 3, \quad p = 4, \quad n_1 = 271, \quad n_2 = 138, \quad n_3 = 107$$

$$\ln |\mathbf{S}_1| = -17.397, \quad \ln |\mathbf{S}_2| = -13.926$$

$$\ln |\mathbf{S}_3| = -15.741, \quad \ln |\mathbf{S}_{pooled}| = -15.564$$

$$u = \left[\frac{1}{270} + \frac{1}{137} + \frac{1}{106} - \frac{1}{270+137+106} \right] \left[\frac{2(4)^2 + 3(4) - 1}{6(4+1)(3-1)} \right]$$

$$= 0.0133$$

$$M = [270 + 137 + 106](-15.564) -$$

$$[270(-17.397) + 137(-13.926) + 106(-15.741)] = 289.3$$

$$C = (1 - 0.0133) \times 289.3 = 285.5$$

$$\nu = 4(4+1)(3-1)/2 = 20$$

$$H_0 \text{ is rejected at any reasonable level of significance from } \chi^2_{\nu} \text{ table for comparison with } C$$

120

Example 6.13: Plastic Film Data

x_1 = tear resistance, x_2 = gloss, and x_3 = opacity

		Factor 2: Amount of additive					
		Low (1.0%)			High (1.5%)		
Factor 1: Change in rate of extrusion	Low (-10%)	x_1	x_2	x_3	x_1	x_2	x_3
		[6.5	9.5	4.4]	[6.9	9.1	5.7]
		[6.2	9.9	6.4]	[7.2	10.0	2.0]
		[5.8	9.6	3.0]	[6.9	9.9	3.9]
	High (10%)	x_1	x_2	x_3	x_1	x_2	x_3
		[6.5	9.6	4.1]	[6.1	9.5	1.9]
		[6.5	9.2	0.8]	[6.3	9.4	5.7]
		[6.7	9.1	2.8]	[7.1	9.2	8.4]

121

Outline

- Comparing Several Multivariate Population Means (One-Way MANOVA)
- Simultaneous Confidence Intervals for Treatment Effects
- Testing for Equality of Covariance Matrices
- Two-Way ANOVA
- Two-Way Multivariate Analysis of Variance

122

Questions

- How to determine if a factor and its interaction with the other factor is significant if two factors are involved in an experiment?
- What are the four types of interactions of two factors?
- What is the two-way ANOVA table?

123

Scenarios

- To observe if effects of factors in the following scenarios are significant
 - Ratings of music compressed by MP3 compressors: brands vs. ages of the subjects
 - Performance of Teaching: methods (Lectures, group studying, and computer assisted instruction) vs. genders of undergraduate students

124

Teaching Methods vs. Gender: Knowing only Overall Mean

Gender	CAI	Lecture	Group Studying	Mean
Boys	50	50	50	50
Girls	50	50	50	50
Mean	50	50	50	50

125

Teaching Methods vs. Gender: Knowing Overall Mean and Row Effects

Gender	CAI	Lecture	Group Studying	Mean
Boys	40	40	40	40
Girls	60	60	60	60
Mean	50	50	50	50

126

Teaching Methods vs. Gender: Knowing Overall Mean, Row Effects, and Column Effects

Gender	CAI	Lecture	Group Studying	Mean
Boys	50	40	30	40
Girls	70	60	50	60
Mean	60	50	40	50

127

Teaching Methods vs. Gender: Including Interaction Terms

Gender	CAI	Lecture	Group Studying	Mean
Boys	65	40	15	40
Girls	55	60	65	60
Mean	60	50	40	50

128

Comparing Four MP3 Compressors

- Test four brands, *A, B, C, D*
- 10 subjects, 5 young and 5 senior, each brand (40 in total) to provide a satisfaction rating on a 10-point scale
- Assume that the rating to each brand is a normal distribution, but all four distributions are with the same variance

129

Sample Data

		A	B	C	D	Mean
Young Subjects	1~4	4	5	7	2	5.05
	5~8	4	5	8	1	
	9~12	5	6	7	2	
	13~16	5	6	9	3	
	17~20	6	7	6	3	
	Mean	4.8	5.8	7.4	2.2	

*Adapted from: G. R. Norman and D. L. Streiner, *Biostatistics*, 3rd ed.¹³⁰

Sample Data

		A	B	C	D	Mean
Senior Subjects	21~24	3	6	3	4	3.70
	25~28	4	4	2	5	
	29~32	4	5	2	4	
	33~36	3	6	2	4	
	37~40	4	3	3	3	
	Mean	3.6	4.8	2.4	4.0	

	A	B	C	D	Mean
Brand Mean	4.2	5.3	4.9	3.1	4.375

*Adapted from: G. R. Norman and D. L. Streiner, *Biostatistics*, 3rd ed.¹³¹

Sum of Squares (Young/Senior)

$$SS(\text{young} / \text{senior}) = bn \sum_{\ell=1}^g (\bar{x}_{\ell \cdot} - \bar{x})^2$$

$$SS(\text{young} / \text{senior}) = 20[(5.05 - 4.375)^2 + (3.70 - 4.375)^2] \\ = 18.225$$

132

Sum of Squares (Brands)

$$SS(\text{brands}) = gn \sum_{k=1}^b (\bar{x}_{\bullet k} - \bar{x})^2$$

$$SS(\text{brands}) = 10[(4.2 - 4.375)^2 + (5.3 - 4.375)^2 + (4.9 - 4.375)^2 + (3.1 - 4.375)^2] = 27.875$$

133

Sum of Squares (Within)

$$SS(\text{within}) = \sum_{\ell=1}^g \sum_{k=1}^b \sum_{\gamma=1}^n (x_{\ell k \gamma} - \bar{x}_{\ell k})^2$$

$$SS(\text{within}) = (4 - 4.8)^2 + (4 - 4.8)^2 + \cdots + (6 - 4.8)^2 + (5 - 5.8)^2 + (5 - 5.8)^2 + \cdots + (7 - 5.9)^2 + \cdots + (4 - 4.0)^2 + (5 - 4.0)^2 + \cdots + (3 - 4.0)^2$$

[40 terms]

$$= 24.80$$

134

Sum of Squares (Total)

$$x_{\ell kr} = \bar{x} + (\bar{x}_{\ell \bullet} - \bar{x}) + (\bar{x}_{\bullet k} - \bar{x}) + (\bar{x}_{\ell k} - \bar{x}_{\ell \bullet} - \bar{x}_{\bullet k} + \bar{x}) + (x_{\ell kr} - \bar{x}_{\ell k})$$

$$\sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (x_{\ell kr} - \bar{x})^2 = \sum_{\ell=1}^g bn(\bar{x}_{\ell \bullet} - \bar{x})^2 + \sum_{k=1}^b gn(\bar{x}_{\bullet k} - \bar{x})^2 + \sum_{\ell=1}^g \sum_{k=1}^b n(\bar{x}_{\ell k} - \bar{x}_{\ell \bullet} - \bar{x}_{\bullet k} + \bar{x})^2 + \sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (x_{\ell kr} - \bar{x}_{\ell k})^2$$

$$SS(\text{total}) = SS(\text{young/senior}) + SS(\text{brand}) + SS(\text{interactions}) + SS(\text{within})$$

135

Sum of Squares (Interactions)

$$SS(\text{interactions}) = n \sum_{\ell=1}^g \sum_{k=1}^b (\bar{x}_{\ell k} - \bar{x}_{\ell \bullet} - \bar{x}_{\bullet k} + \bar{x})^2$$

$$SS(\text{interactions}) = 5[(4.8 - 4.875)^2 + (3.6 - 3.525)^2 + \cdots + (4.0 - 2.425)^2]$$

[8 terms]

$$= 58.475$$

136

Sum of Squares (Total)

$$\begin{aligned}
 SS(total) &= (4 - 4.375)^2 + (4 - 4.375)^2 + \dots + (4 - 4.375)^2 + \\
 &\quad (5 - 4.375)^2 + (5 - 4.375)^2 + \dots + (3 - 4.375)^2 + \\
 &\quad (7 - 4.375)^2 + (8 - 4.375)^2 + \dots + (3 - 4.375)^2 + \\
 &\quad (2 - 4.375)^2 + (1 - 4.375)^2 + \dots + (3 - 4.375)^2 \\
 &\quad [40 \text{ terms}] \\
 &= 129.375 = 18.225 + 58.475 + 24.80 + 27.875
 \end{aligned}$$

137

Expected Values of Sum of Squares

$$E[SS(\text{brand}) / df(\text{brand})] \text{ contains } \sigma_{\text{brand}}^2, \sigma_{\text{interactions}}^2, \sigma_{\text{err}}^2$$

$$E[SS(\text{within}) / df(\text{within})] = \sigma_{\text{err}}^2$$

Thus, if brand effect is significant

$$\frac{E[SS(\text{brand}) / df(\text{brand})]}{E[SS(\text{within}) / df(\text{within})]} > 1$$

$$F_{\text{brand}} > 1$$

138

Degrees of Freedom

$$df(\text{young} / \text{senior}) = g - 1 = 2 - 1 = 1$$

$$df(\text{brand}) = b - 1 = 4 - 1 = 3$$

$$df(\text{within}) = bg(n - 1) = 8(5 - 1) = 32$$

$$df(\text{interactions}) = (b - 1)(g - 1) = (4 - 1)(2 - 1) = 3$$

$$\begin{aligned}
 df(\text{total}) &= bgn - 1 = bg(n - 1) + (b - 1)(g - 1) + b - 1 + g - 1 \\
 &= df(\text{within}) + df(\text{interactions}) + \\
 &\quad df(\text{brand}) + df(\text{young} / \text{senior}) \\
 &= 40 - 1 = 39 = 32 + 3 + 3 + 1
 \end{aligned}$$

139

Two-way ANOVA Summary

Source	Sum of Squares	df	Mean square	F
Brand	27.875	3	9.29	11.99
Young/ Senior	18.225	1	18.23	23.52
Brand X Y/S	58.475	3	19.49	25.15
Within	24.80	32	0.78	
Total	129.375	39		

140

Hypothesis Testing

$$F_{brand} = 11.99 > F_{3,32}(0.05) \approx 2.92$$


$$F_{Y/S} = 23.52 > F_{1,32}(0.05) \approx 4.17$$

$$F_{interactions} = 25.15 > F_{3,32}(0.05) \approx 2.92$$

All factors and interactions are significant at 0.05 significance level


141

Histogram of Means




142

Effect of Interaction



143

Possible Types of Interactions




144

de Groot's Experiment (1965)

- ✦ Observed the ability of chess masters and novices to recall piece positions
- ✦ Experts
 - Recalled about 90% of the pieces in a typical mid-game
- ✦ Novices
 - Recalled about 20%
- ✦ Many factors might have been introduced
- ✦ Randomized piece positions
 - Everybody recalled about 20%
 - No effect of expertise

145

Interpretation by Adjusted Data



146

Two-Way ANOVA

$$X_{\ell kr} = \mu + \tau_{\ell} + \beta_k + \gamma_{\ell k} + e_{\ell kr}$$

$$\ell = 1, 2, \dots, g; \quad k = 1, 2, \dots, b; \quad r = 1, 2, \dots, n$$

$$\sum_{\ell=1}^g \tau_{\ell} = \sum_{k=1}^b \beta_k = \sum_{\ell=1}^g \gamma_{\ell k} = \sum_{k=1}^b \gamma_{\ell k} = 0, \quad e_{\ell kr} : N(0, \sigma^2)$$

$$E(X_{\ell kr}) = \mu + \tau_{\ell} + \beta_k + \gamma_{\ell k}$$

$$x_{\ell kr} = \bar{x} + (\bar{x}_{\ell \cdot} - \bar{x}) + (\bar{x}_{\cdot k} - \bar{x}) + (\bar{x}_{\ell k} - \bar{x}_{\ell \cdot} - \bar{x}_{\cdot k} + \bar{x}) + (x_{\ell kr} - \bar{x}_{\ell k})$$


$$\sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (x_{\ell kr} - \bar{x})^2 = \sum_{\ell=1}^g bn(\bar{x}_{\ell \cdot} - \bar{x})^2 + \sum_{k=1}^b gn(\bar{x}_{\cdot k} - \bar{x})^2$$

$$+ \sum_{\ell=1}^g \sum_{k=1}^b n(\bar{x}_{\ell k} - \bar{x}_{\ell \cdot} - \bar{x}_{\cdot k} + \bar{x})^2 + \sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (x_{\ell kr} - \bar{x}_{\ell k})^2$$

$$SS_{cor} = SS_{fac1} + SS_{fac2} + SS_{int} + SS_{res}$$

147

Effect of Interactions



148

Two-Way ANOVA

ANOVA TABLE FOR COMPARING EFFECTS OF TWO FACTORS AND THEIR INTERACTION

Source of variation	Sum of squares (SS)	Degrees of freedom (d.f.)
Factor 1	$SS_{fac1} = \sum_{\ell=1}^g bn(\bar{x}_{\ell.} - \bar{x})^2$	$g - 1$
Factor 2	$SS_{fac2} = \sum_{k=1}^b gn(\bar{x}_{.k} - \bar{x})^2$	$b - 1$
Interaction	$SS_{int} = \sum_{\ell=1}^g \sum_{k=1}^b n(\bar{x}_{\ell k} - \bar{x}_{\ell.} - \bar{x}_{.k} + \bar{x})^2$	$(g - 1)(b - 1)$
Residual (Error)	$SS_{res} = \sum_{\ell=1}^g \sum_{k=1}^b \sum_{i=1}^n (x_{\ell ki} - \bar{x}_{\ell k})^2$	$gb(n - 1)$
Total (corrected)	$SS_{cor} = \sum_{\ell=1}^g \sum_{k=1}^b \sum_{i=1}^n (x_{\ell ki} - \bar{x})^2$	$gbn - 1$

149

Two-Way ANOVA

F - ratio tests

$$\frac{SS_{fac1} / (g - 1)}{SS_{res} / (gb(n - 1))} : \text{for effects of factor 1}$$

$$\frac{SS_{fac2} / (b - 1)}{SS_{res} / (gb(n - 1))} : \text{for effects of factor 2}$$

$$\frac{SS_{int} / (g - 1)(b - 1)}{SS_{res} / (gb(n - 1))} : \text{for effects of factor 1 - factor 2 interaction}$$

150

Outline

- Comparing Several Multivariate Population Means (One-Way MANOVA)
- Simultaneous Confidence Intervals for Treatment Effects
- Testing for Equality of Covariance Matrices
- Two-Way ANOVA
- Two-Way Multivariate Analysis of Variance

151

Questions

- What is the two-way MANOVA table?
- How to determine if the interaction effect exists?
- How to test the effect of each factor by the two-way MANOVA?
- How to determine the Bonferroni confidence intervals if the interaction effect is negligible?

152

Two-Way MANOVA

$$\mathbf{X}_{\ell kr} = \boldsymbol{\mu} + \boldsymbol{\tau}_{\ell} + \boldsymbol{\beta}_k + \boldsymbol{\gamma}_{\ell k} + \mathbf{e}_{\ell kr}$$

$$\ell = 1, 2, \dots, g; \quad k = 1, 2, \dots, b; \quad r = 1, 2, \dots, n$$

$$\sum_{\ell=1}^g \boldsymbol{\tau}_{\ell} = \sum_{k=1}^b \boldsymbol{\beta}_k = \sum_{\ell=1}^g \boldsymbol{\gamma}_{\ell k} = \sum_{k=1}^b \boldsymbol{\gamma}_{\ell k} = \mathbf{0}, \quad \mathbf{e}_{\ell kr} : N_p(\mathbf{0}, \boldsymbol{\Sigma})$$

$$\mathbf{x}_{\ell kr} = \bar{\mathbf{x}} + (\bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}}) + (\bar{\mathbf{x}}_{\cdot k} - \bar{\mathbf{x}}) + (\bar{\mathbf{x}}_{\ell k} - \bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}}_{\cdot k} + \bar{\mathbf{x}}) + (\mathbf{x}_{\ell kr} - \bar{\mathbf{x}}_{\ell k})$$

$$\begin{aligned} \sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (\mathbf{x}_{\ell kr} - \bar{\mathbf{x}})(\mathbf{x}_{\ell kr} - \bar{\mathbf{x}})' = & \sum_{\ell=1}^g bn(\bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}})' + \sum_{k=1}^b gn(\bar{\mathbf{x}}_{\cdot k} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\cdot k} - \bar{\mathbf{x}})' \\ & + \sum_{\ell=1}^g \sum_{k=1}^b n(\bar{\mathbf{x}}_{\ell k} - \bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}}_{\cdot k} + \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell k} - \bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}}_{\cdot k} + \bar{\mathbf{x}})' \\ & + \sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (\mathbf{x}_{\ell kr} - \bar{\mathbf{x}}_{\ell k})(\mathbf{x}_{\ell kr} - \bar{\mathbf{x}}_{\ell k})' \end{aligned}$$

153

Two-Way MANOVA

MANOVA TABLE FOR COMPARING FACTORS AND THEIR INTERACTION

Source of variation	Matrix of sum of squares and cross products (SSP)	Degrees of freedom (d.f.)
Factor 1	$SSP_{fac1} = \sum_{\ell=1}^g bn(\bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}})'$	$g - 1$
Factor 2	$SSP_{fac2} = \sum_{k=1}^b gn(\bar{\mathbf{x}}_{\cdot k} - \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\cdot k} - \bar{\mathbf{x}})'$	$b - 1$
Interaction	$SSP_{int} = \sum_{\ell=1}^g \sum_{k=1}^b n(\bar{\mathbf{x}}_{\ell k} - \bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}}_{\cdot k} + \bar{\mathbf{x}})(\bar{\mathbf{x}}_{\ell k} - \bar{\mathbf{x}}_{\ell \cdot} - \bar{\mathbf{x}}_{\cdot k} + \bar{\mathbf{x}})'$	$(g - 1)(b - 1)$
Residual (Error)	$SSP_{res} = \sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (\mathbf{x}_{\ell kr} - \bar{\mathbf{x}}_{\ell k})(\mathbf{x}_{\ell kr} - \bar{\mathbf{x}}_{\ell k})'$	$gb(n - 1)$
Total (corrected)	$SSP_{cor} = \sum_{\ell=1}^g \sum_{k=1}^b \sum_{r=1}^n (\mathbf{x}_{\ell kr} - \bar{\mathbf{x}})(\mathbf{x}_{\ell kr} - \bar{\mathbf{x}})'$	$gbn - 1$

154

Two-Way MANOVA

Test for interaction

For large samples, reject $H_0 : \boldsymbol{\gamma}_{11} = \boldsymbol{\gamma}_{12} = \dots = \boldsymbol{\gamma}_{gb} = \mathbf{0}$ if

$$-\left[gb(n-1) - \frac{p+1-(g-1)(b-1)}{2} \right] \ln \Lambda^* > \chi^2_{(g-1)(b-1)}(\alpha)$$

$$\text{Wilk's lambda } \Lambda^* = \frac{|\text{SSP}_{res}|}{|\text{SSP}_{int} + \text{SSP}_{res}|}$$

If interaction effects exist, the factor effects do not have a clear interpretation

155

Two-Way MANOVA

Test for factor 1 effect

For large samples, reject $H_0 : \boldsymbol{\tau}_1 = \boldsymbol{\tau}_2 = \dots = \boldsymbol{\tau}_g = \mathbf{0}$ if

$$-\left[gb(n-1) - \frac{p+1-(g-1)(b-1)}{2} \right] \ln \Lambda^* > \chi^2_{(g-1)p}(\alpha)$$

$$\text{Wilk's lambda } \Lambda^* = \frac{|\text{SSP}_{res}|}{|\text{SSP}_{fac1} + \text{SSP}_{res}|}$$

Test for factor 2 effect

For large samples, reject $H_0 : \boldsymbol{\beta}_1 = \boldsymbol{\beta}_2 = \dots = \boldsymbol{\beta}_b = \mathbf{0}$ if

$$-\left[gb(n-1) - \frac{p+1-(g-1)(b-1)}{2} \right] \ln \Lambda^* > \chi^2_{(b-1)p}(\alpha)$$

$$\text{Wilk's lambda } \Lambda^* = \frac{|\text{SSP}_{res}|}{|\text{SSP}_{fac2} + \text{SSP}_{res}|}$$

156

Bonferroni Confidence Intervals

With negligible interactions,
the simultaneous confidence intervals are

$$(\bar{x}_{\ell \cdot i} - \bar{x}_{m \cdot i}) \pm t_p \left(\frac{\alpha}{pg(g-1)} \right) \sqrt{\frac{E_{ii}}{\nu} \frac{2}{bn}} \quad \text{for } \tau_{\ell i} - \tau_{mi}$$

and

$$(\bar{x}_{\cdot ki} - \bar{x}_{\cdot qi}) \pm t_p \left(\frac{\alpha}{pb(b-1)} \right) \sqrt{\frac{E_{ii}}{\nu} \frac{2}{gn}} \quad \text{for } \beta_{ki} - \beta_{qi}$$

$$\nu = gb(n-1), \quad \mathbf{E} = \text{SSP}_{res}$$

157

Example 6.13: MANOVA Table

Source of variation	SSP			d.f.
Factor 1: change in rate of extrusion	1.7405	-1.5045	.8555	1
		1.3005	-.7395	
Factor 2: amount of additive	.7605	.6825	1.9305	1
		.6125	1.7325	
Interaction	.0005	.0165	.0445	1
		.5445	1.4685	
Residual	1.7640	.0200	-3.0700	16
		2.6280	-.5520	
Total (corrected)	4.2655	-.7855	-.2395	19
		5.0855	1.9095	
			74.2055	

158

Example 6.13: Interaction

$$\Lambda^* = \frac{|\text{SSP}_{res}|}{|\text{SSP}_{int} + \text{SSP}_{res}|} = 0.7771$$

$$(g-1)(b-1) = 1$$

$$F = \left(\frac{1 - \Lambda^*}{\Lambda^*} \right) \frac{(gb(n-1) - p + 1)/2}{((g-1)(b-1) - p + 1)/2} : F_{\nu_1, \nu_2}$$

$$\nu_1 = |(g-1)(b-1) - p| + 1 = 3$$

$$\nu_2 = gb(n-1) - p + 1 = 14$$

$$F = 1.34 < F_{3,14}(0.05) = 3.34$$

$$H_0 : \gamma_{11} = \gamma_{12} = \gamma_{21} = \gamma_{22} = 0 \text{ (no interaction) is not rejected}$$

159

Example 6.13: Effects of Factors 1 & 2

$$\Lambda_1^* = \frac{|\text{SSP}_{res}|}{|\text{SSP}_{fac1} + \text{SSP}_{res}|} = 0.3819$$

$$\Lambda_2^* = \frac{|\text{SSP}_{res}|}{|\text{SSP}_{fac2} + \text{SSP}_{res}|} = 0.5230$$

$$F_1 = \left(\frac{1 - \Lambda_1^*}{\Lambda_1^*} \right) \frac{\nu_2/2}{\nu_1/2} = 7.55, \quad \nu_1 = |(g-1) - p| + 1 = 3$$

$$F_2 = \left(\frac{1 - \Lambda_2^*}{\Lambda_2^*} \right) \frac{\nu_2/2}{\nu_1/2} = 4.26, \quad \nu_1 = |(b-1) - p| + 1 = 3$$

$$\nu_2 = gb(n-1) - p + 1 = 14$$

$$F_1 > F_{3,14}(0.05) = 3.34, \quad \text{reject } H_0 : \tau_1 = \tau_2 = 0$$

$$F_2 > F_{3,14}(0.05) = 3.34, \quad \text{reject } H_0 : \beta_1 = \beta_2 = 0$$

160

Outline

- Profile Analysis
- ANOVA for Repeated Measures
- Repeated Measures Designs and Growth Curves
- Perspectives and Strategy for Analyzing Multivariate Models

161

Questions

- What is the profile analysis?
- How to carry out the profile analysis?


162

Profile Analysis

- A battery of p treatments (tests, questions, etc.) are administered to two or more group of subjects
- The question of equality of mean vectors is divided into several specific possibilities
 - Are the profiles parallel?
 - Are the profiles coincident?
 - Are the profiles level?


163

Example 6.14: Love and Marriage Data



164

Population Profile



165

Profile Analysis

Assume two populations

Are the profiles parallel?

$$H_{01} : \mu_{1i} - \mu_{1i-1} = \mu_{2i} - \mu_{2i-1}, i = 2, 3, \dots, p$$

Are the profiles coincident?

$$H_{02} : \mu_{1i} = \mu_{2i}, i = 1, 2, \dots, p$$

Are the profiles level?

$$H_{03} : \mu_{11} = \mu_{12} = \dots = \mu_{1p} = \mu_{21} = \mu_{22} = \dots = \mu_{2p}$$

166

Test for Parallel Profiles

$$\mathbf{C}_{(p-1) \times p} = \begin{bmatrix} -1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & -1 & 1 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & -1 & 1 \end{bmatrix}$$

$$\mathbf{C}\mathbf{X}_{1j} : N_{p-1}(\mathbf{C}\boldsymbol{\mu}_1, \mathbf{C}\boldsymbol{\Sigma}\boldsymbol{\Sigma}'), \mathbf{C}\mathbf{X}_{2j} : N_{p-1}(\mathbf{C}\boldsymbol{\mu}_2, \mathbf{C}\boldsymbol{\Sigma}\boldsymbol{\Sigma}')$$

Reject $H_{01} : \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{C}\boldsymbol{\mu}_2$ at level α if

$$T^2 = (\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2)' \mathbf{C}' \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{C}\mathbf{S}_{pooled} \mathbf{C}' \right]^{-1} \mathbf{C}(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2) > c^2$$

$$c^2 = \frac{(n_1 + n_2 - 2)(p - 1)}{n_1 + n_2 - p} F_{p-1, n_1 + n_2 - p}(\alpha)$$

167

Test for Coincident Profiles

Given parallel profiles

Reject $H_{02} : \mathbf{1}'\boldsymbol{\mu}_1 = \mathbf{1}'\boldsymbol{\mu}_2$ at level α if

$$T^2 = \mathbf{1}'(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2) \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{1}'\mathbf{S}_{pooled}\mathbf{1} \right]^{-1} \mathbf{1}'(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2)$$

$$= \left(\frac{\mathbf{1}'(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2)}{\sqrt{\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{1}'\mathbf{S}_{pooled}\mathbf{1}}} \right)^2 > t_{n_1 + n_2 - 2}^2 \left(\frac{\alpha}{2} \right) = F_{1, n_1 + n_2 - 2}(\alpha)$$

168

Test for Level Profiles

Given coincident profiles

Reject $H_{03} : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ at level α if

$$(n_1 + n_2) \bar{\mathbf{x}}' \mathbf{C}' [\mathbf{CSC}']^{-1} \mathbf{C} \bar{\mathbf{x}} > c^2$$

$$c^2 = \frac{(n_1 + n_2 - 1)(p - 1)}{n_1 + n_2 - p - 1} F_{p-1, n_1 + n_2 - p + 1}(\alpha)$$

$$\bar{\mathbf{x}} = \frac{n_1}{n_1 + n_2} \bar{\mathbf{x}}_1 + \frac{n_2}{n_1 + n_2} \bar{\mathbf{x}}_2$$

169

Example 6.14

$$\bar{\mathbf{x}}_1 = \begin{bmatrix} 6.833 \\ 7.033 \\ 3.967 \\ 4.700 \end{bmatrix}, \bar{\mathbf{x}}_2 = \begin{bmatrix} 6.633 \\ 7.000 \\ 4.000 \\ 4.533 \end{bmatrix}$$

$$\mathbf{S}_{pooled} = \begin{bmatrix} 0.606 & 0.262 & 0.066 & 0.161 \\ 0.262 & 0.637 & 0.173 & 0.143 \\ 0.066 & 0.173 & 0.810 & 0.029 \\ 0.161 & 0.143 & 0.029 & 0.306 \end{bmatrix}$$

170

Example 6.14: Test for Parallel Profiles


$$\mathbf{CS}_{pooled} \mathbf{C}' = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \mathbf{S}_{pooled} \begin{bmatrix} -1 & 0 & 0 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\mathbf{C}(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2) = \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 0.200 \\ 0.033 \\ -0.033 \\ 0.167 \end{bmatrix} = \begin{bmatrix} -0.167 \\ -0.066 \\ 0.200 \end{bmatrix}$$

$$T^2 = 1.005 < \frac{(30 + 30 - 2)(4 - 1)}{30 + 30 - 4} F_{3,56}(0.05) = 8.7$$

171

Example 6.14: Sample Profiles



172

Example 6.14: Test for Coincident Profiles

$$\mathbf{1}'(\bar{\mathbf{x}}_1 - \bar{\mathbf{x}}_2) = 0.367$$

$$\mathbf{1}'\mathbf{S}_{pooled}\mathbf{1} = 4.207$$

$$T^2 = \left(\frac{0.367}{\sqrt{\left(\frac{1}{30} + \frac{1}{30}\right)4.207}} \right)^2 = 0.501 < F_{1,58}(0.05) = 4.0$$

173

Outline

- Profile Analysis
- ANOVA for Repeated Measures
- Repeated Measures Designs and Growth Curves
- Perspectives and Strategy for Analyzing Multivariate Models

174

Questions

- What are repeated measures?
- How to view the data for repeated measures in a two-way ANOVA view?
- How to test the null hypothesis in repeated measures?

175

Repeated-Measures ANOVA

- Drugs A, B, C are tested to see if they are equally effective for pain relief
- Subjects are to take all of the drugs, in turn, suitably blinded and after a suitable washout period
- Subjects rate the degree of pain belief on a 1 to 6 scale (1: no relief, 6 complete relief)

176

Avoiding Order Effects

- ✦ Randomize the order of treatment
 - 1/3 get drug A first, 1/3 get drug B first, 1/3 get drug C first
- ✦ People in a long, natural healing course may grow tolerant of the irritant and learn to tune them out
 - The last medication may work the best
 - Order effects

177

Sample Data

Subject	A	B	C	Average
1	5	3	2	3.33
2	5	4	3	4.00
3	5	6	5	5.33
4	6	4	2	4.00
5	6	6	6	6.00
6	4	2	1	2.33
7	4	4	3	3.67
8	4	5	5	4.67
9	4	2	2	2.67
10	5	3	1	3.00
Means	4.80	3.90	3.00	3.90

*Adapted from: G. R. Norman and D. L. Streiner, *Biostatistics*, 3rd ed.¹⁷⁸

Two-Way ANOVA View

- ✦ Individual subjects as one factor
- ✦ Pain reliever as a second factor
- ✦ Cells are defined by
 - Subjects: 10 levels
 - Drug: 3 levels
- ✦ One observation per cell
- ✦ Special case of two-way ANOVA
 - $n = 1, g = 10, b = 3$

179

Sum of Squares (Drug)

$$SS(\text{drug}) = g \sum_{k=1}^b (\bar{x}_{\bullet k} - \bar{x})^2$$

$$SS(\text{drug}) = 10[(4.8 - 3.9)^2 + (3.9 - 3.9)^2 + (3.0 - 3.9)^2] \\ = 16.2$$

180

Sum of Squares (Subjects)

$$SS(subjects) = b \sum_{\ell=1}^g (\bar{x}_{\ell\bullet} - \bar{x})^2$$

$$SS(subjects) = 3[(3.33 - 3.90)^2 + (4.00 - 3.90)^2 + \dots + (3.00 - 3.90)^2] \\ = 36.7$$

181

Sum of Squares (Interaction)

$$SS(interaction) = \sum_{\ell=1}^g \sum_{k=1}^b (\bar{x}_{\ell k} - \bar{x}_{\ell\bullet} - \bar{x}_{\bullet k} + \bar{x})^2$$

$$SS(interaction) = [(5 - 4.23)^2 + (3 - 3.33)^2 + \dots + (1 - 2.10)^2] \\ [30 \text{ terms}] \\ = 15.8$$

182

Sum of Squares (Within)

$$SS(within) = \sum_{\ell=1}^g \sum_{k=1}^b (x_{\ell k\gamma} - \bar{x}_{\ell k})^2 = 0$$

183

Degrees of Freedom

$$df(subject) = g - 1 = 10 - 1 = 9$$

$$df(drug) = b - 1 = 3 - 1 = 2$$

$$df(within) = bg(n - 1) = 0$$

$$df(interaction) = (b - 1)(g - 1) = (3 - 1)(10 - 1) = 18$$

$$df(total) = bg - 1 = (b - 1)(g - 1) + b - 1 + g - 1$$

$$= df(interaction) +$$

$$df(drug) + df(subject)$$

$$= 30 - 1 = 29 = 18 + 2 + 9$$

184

Signal vs. Noise

- ✦ To determine if there is any significant difference in relief from different pain relievers
 - Main effect of Drug
- ✦ $SS(within) = 0$
- ✦ Choose $SS(interaction)$ as error term
 - Reflects the extent to which different subjects respond differently to the different drug types

185

ANOVA Table

Source	Sum of Squares	<i>df</i>	Mean square	F
Drug	16.2	2	8.100	9.225
Subject	36.7	9	4.078	
Drug X Subject	15.8	18	0.878	
Totals	68.7	29		

186

Hypothesis Testing

$$F_{Drug} = 9.225 > F_{2,18}(0.05) \approx 3.55$$

Drug effect is significant (i.e., difference exists)
at 0.05 significance level

187

ANOVA Table for Same Data as a One-Way ANOVA Test

Source	Sum of Squares	<i>df</i>	Mean square	F
Drug	16.2	2	8.100	4.107
Error	52.5	27	1.944	
Totals	68.7	29		

188

Outline

- Profile Analysis
- ANOVA for Repeated Measured
- Repeated Measures Designs and Growth Curves
- Perspectives and Strategy for Analyzing Multivariate Models

189

Questions

- How to compare growth curves?

190

Example 6.15: Ulna Data, Control Group

Subject	Initial	1 year	2 year	3 year
1	87.3	86.9	86.7	75.5
2	59.0	60.2	60.0	53.6
3	76.7	76.5	75.7	69.5
4	70.6	76.1	72.1	65.3
5	54.9	55.1	57.2	49.0
6	78.2	75.3	69.1	67.6
7	73.7	70.8	71.8	74.6
8	61.8	68.7	68.2	57.4
9	85.3	84.4	79.2	67.0
10	82.3	86.9	79.4	77.4
11	68.6	65.4	72.3	60.8
12	67.8	69.2	66.3	57.9
13	66.2	67.0	67.0	56.2
14	81.0	82.3	86.8	73.9
15	72.3	74.6	75.3	66.1
Mean	72.38	73.29	72.47	64.79

Source: Data courtesy of Everett Smith.

191

Example 6.15: Ulna Data, Treatment Group

Subject	Initial	1 year	2 year	3 year
1	83.8	85.5	86.2	81.2
2	65.3	66.9	67.0	60.6
3	81.2	79.5	84.5	75.2
4	75.4	76.7	74.3	66.7
5	55.3	58.3	59.1	54.2
6	70.3	72.3	70.6	68.6
7	76.5	79.9	80.4	71.6
8	66.0	70.9	70.3	64.1
9	76.7	79.0	76.9	70.3
10	77.2	74.0	77.8	67.9
11	67.3	70.7	68.9	65.9
12	50.3	51.4	53.6	48.0
13	57.7	57.0	57.5	51.5
14	74.3	77.7	72.6	68.0
15	74.0	74.7	74.5	65.7
16	57.3	56.0	64.7	53.0
Mean	69.29	70.66	71.18	64.53

Source: Data courtesy of Everett Smith.

192

Comparison of Growth Curves

\mathbf{X}_{ij} : vector of p measurements on subject j in group ℓ

$j = 1, 2, \dots, n_\ell$; $\ell = 1, 2, \dots, g$

\mathbf{X}_{ij} : Multivariate normal with covariance Σ

Putthoff - Roy model

$$E(\mathbf{X}_{ij}) = \begin{bmatrix} \beta_{i0} + \beta_{i1}t_1 + \dots + \beta_{iq}t_1^q \\ \beta_{i0} + \beta_{i1}t_2 + \dots + \beta_{iq}t_2^q \\ \vdots \\ \beta_{i0} + \beta_{i1}t_p + \dots + \beta_{iq}t_p^q \end{bmatrix} = \begin{bmatrix} 1 & t_1 & \dots & t_1^q \\ 1 & t_2 & \dots & t_2^q \\ \vdots & \vdots & \ddots & \vdots \\ 1 & t_p & \dots & t_p^q \end{bmatrix} \begin{bmatrix} \beta_{i0} \\ \beta_{i1} \\ \vdots \\ \beta_{iq} \end{bmatrix} \\ = \mathbf{B}\boldsymbol{\beta}_\ell$$

193

Comparison of Growth Curves

Maximum likelihood estimators of $\boldsymbol{\beta}_\ell$:

$$\hat{\boldsymbol{\beta}}_\ell = (\mathbf{B}'\mathbf{S}_{pooled}^{-1}\mathbf{B})^{-1}\mathbf{B}'\mathbf{S}_{pooled}^{-1}\bar{\mathbf{X}}_\ell$$

$$\mathbf{S}_{pooled} = \frac{1}{N-g}((n_1-1)\mathbf{S}_1 + \dots + (n_g-1)\mathbf{S}_g) = \frac{\mathbf{W}}{N-g}$$

$$N = \sum_{\ell=1}^g n_\ell, \quad \hat{\text{Cov}}(\hat{\boldsymbol{\beta}}_\ell) = \frac{k}{n_\ell}(\mathbf{B}'\mathbf{S}_{pooled}^{-1}\mathbf{B})^{-1}$$

$$k = (N-g)(N-g-1)/(N-g-p+q)(N-g-p+q+1)$$

$$\mathbf{W}_q = \sum_{\ell=1}^g \sum_{j=1}^{n_\ell} (\mathbf{X}_{ij} - \mathbf{B}\hat{\boldsymbol{\beta}}_\ell)(\mathbf{X}_{ij} - \mathbf{B}\hat{\boldsymbol{\beta}}_\ell)', \quad \Lambda^* = \frac{|\mathbf{W}|}{|\mathbf{W}_q|}$$

Reject the null hypothesis that the polynomial is adequate if

$$-(N-(p-q+g)/2)\ln \Lambda^* > \chi_{(p-q-1)g}^2(\alpha)$$

194

Example 6.15

Use quadratic growth model

$$\begin{bmatrix} \hat{\beta}_1 & \hat{\beta}_2 \end{bmatrix} = \begin{bmatrix} 73.0701 (2.58) & 70.1387 (2.50) \\ 3.6444 (0.83) & 4.0900 (0.80) \\ -2.0274 (0.28) & -1.8534 (0.27) \end{bmatrix}$$

Control Group: $73.07 + 3.64t - 2.03t^2$

Treatment Group: $70.14 + 4.09t - 1.85t^2$

$$\Lambda^* = 0.7627$$

$$-(N-(p-q+g)/2)\ln \Lambda^* = 7.86 < \chi_{(4-2-1)2}^2(0.01) = 9.21$$

195

Outline

- Profile Analysis
- ANOVA for Repeated Measured
- Repeated Measures Designs and Growth Curves
- Perspectives and Strategy for Analyzing Multivariate Models


196

Questions

- What are the strategies in multivariate analysis?
- Why is the experimental design important?

197

Example 6.16: Comparing Multivariate and Univariate Tests



198

Example 6.16: Comparing Multivariate and Univariate Tests

Univariate test on x_1 : $F = 2.46 < F_{1,18}(0.10) = 3.01$

Univariate test on x_2 : $F = 2.68 < F_{1,18}(0.10) = 3.01$

Accept $\mu_1 = \mu_2$

Hotelling's test :

$$T^2 = 17.29 > c^2 = \frac{18 \times 2}{17} F_{2,17}(0.01) = 12.94$$

Reject $\mu_1 = \mu_2$

199

Strategy for Multivariate Comparison of Treatments

- Try to identify outliers
 - Perform calculations with and without the outliers
- Perform a multivariate test of hypothesis
- Calculate the Bonferroni simultaneous confidence intervals
 - For all pairs of groups or treatments, and all characteristics

200

Importance of Experimental Design

- Differences could appear in only one of the many characteristics or a few treatment combinations
- Differences may become lost among all the inactive ones
- Best preventative is a good experimental design
 - Do not include too many other variables that are not expected to show differences

201